# Product Name: Phospho-JAK2 (Y1007 + Y1008) (17F11) Enkilife

**Rabbit Monoclonal Antibody** 

Catalog #: AMRe05930



# Summary

Phospho-JAK2 (Y1007 + Y1008) (17F11) Rabbit Monoclonal Antibody **Production Name** 

Description Rabbit Monoclonal Antibody

Host Rabbit

**Application** WB,IHC-P,ICC/IF,FC,IF-P

Reactivity Human, Mouse, Rat

### **Performance**

Conjugation Unconjugated

Phospho Antibody Modification

Isotype IqG

Clonality Monoclonal

Liquid **Form** 

Storage Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.

Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% New type

**Buffer** preservative N and 50% glycerol. Store at +4°C short term. Store at -20°C long term.

Avoid freeze / thaw cycle.

Affinity purification **Purification** 

### **Immunogen**

**Gene Name** JAK2

**Alternative Names** EC 2.7.10.2; JAK-2; JAK2; Janus kinase 2; kinase Jak2;

Gene ID 3717.0 SwissProt ID O60674.


# **Application**

**Dilution Ratio** WB 1:1000-1:15000, IHC-P/IF-P 1:200-1:1000, ICC/IF 1:200-1:1000, FCM 1:20

**Molecular Weight** 131kDa

Web: https://www.enkilife.com E-mail: order@enkilife.com techsupport@enkilife.com Tel: 0086-27-87002838

Catalog #: AMRe05930




# Background


This gene product is a protein tyrosine kinase involved in a specific subset of cytokine receptor signaling pathways. It has been found to be constitutively associated with the prolactin receptor and is required for responses to gamma interferon. Non-receptor tyrosine kinase involved in various processes such as cell growth, development, differentiation or histone modifications. Mediates essential signaling events in both innate and adaptive immunity. In the cytoplasm, plays a pivotal role in signal transduction via its association with type I receptors such as growth hormone (GHR), prolactin (PRLR), leptin (LEPR), erythropoietin (EPOR), thrombopoietin (THPO); or type II receptors including IFN-alpha, IFN- beta, IFN-gamma and multiple interleukins (PubMed:<a href="http://www.uniprot.org/citations/7615558" target=" blank">7615558</a>). Following ligand-binding to cell surface receptors, phosphorylates specific tyrosine residues on the cytoplasmic tails of the receptor, creating docking sites for STATs proteins (PubMed:<a href="http://www.uniprot.org/citations/9618263" target=" blank">9618263</a>). Subsequently, phosphorylates the STATs proteins once they are recruited to the receptor. Phosphorylated STATs then form homodimer or heterodimers and translocate to the nucleus to activate gene transcription. For example, cell stimulation with erythropoietin (EPO) during erythropoiesis leads to JAK2 autophosphorylation, activation, and its association with erythropoietin receptor (EPOR) that becomes phosphorylated in its cytoplasmic domain. Then, STAT5 (STAT5A or STAT5B) is recruited, phosphorylated and activated by JAK2. Once activated, dimerized STAT5 translocates into the nucleus and promotes the transcription of several essential genes involved in the modulation of erythropoiesis. Part of a signaling cascade that is activated by increased cellular retinol and that leads to the activation of STAT5 (STAT5A or STAT5B) (PubMed: <a href="http://www.uniprot.org/citations/21368206" target=" blank">21368206</a>). In addition, JAK2 mediates angiotensin-2-induced ARHGEF1 phosphorylation (PubMed: <a href="http://www.uniprot.org/citations/20098430" target=" blank">20098430</a>). Plays a role in cell cycle by phosphorylating CDKN1B (PubMed: <a href="http://www.uniprot.org/citations/21423214" target=" blank">21423214</a>). Cooperates with TEC through reciprocal phosphorylation to mediate cytokine-driven activation of FOS transcription. In the nucleus, plays a key role in chromatin by specifically mediating phosphorylation of 'Tyr-41' of histone H3 (H3Y41ph), a specific tag that promotes exclusion of CBX5 (HP1 alpha) from chromatin (PubMed:<a href="http://www.uniprot.org/citations/19783980" target=" blank">19783980</a>).

### Research Area

# **Image Data**

Web: https://www.enkilife.com E-mail: order@enkilife.com techsupport@enkilife.com Tel: 0086-27-87002838





Western blot analysis of JAK2 phosphorylation expression in Jurkat cell lysates treated with Pervanadate.

### Note

For research use only.