

Summary

Production Name	Apolipoprotein CI (10F18) Rabbit Monoclonal Antibody	
Description	Rabbit Monoclonal Antibody	
Host	Rabbit	
Application	WB,IHC-P,IP	
Reactivity	Human	

Performance

Conjugation	Unconjugated
Modification	Unmodified
lsotype	lgG
Clonality	Monoclonal
Form	Liquid
Storage	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.
	Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% New type
Buffer	preservative N and 50% glycerol. Store at +4°C short term. Store at -20°C long term.
	Avoid freeze / thaw cycle.
Purification	Affinity purification

Immunogen

Gene Name	APOC1
Alternative Names	APO C1; Apo Cl; APOC 1; ApoC l; APOC1; APOC1B; ApolipoproteinCl;
Gene ID	341.0
SwissProt ID	P02654.

Application

Dilution Ratio	WB 1:1000, IHC-P/IF-P 1:200-1:1000, IP 1:20-1:50
Molecular Weight	9kDa



Background

Appears to modulate the interaction of APOE with beta-migrating VLDL and inhibit binding of beta-VLDL to the LDL receptor-related protein. Binds free fatty acids and reduces their intracellular esterification. Inhibitor of lipoprotein binding to the low density lipoprotein (LDL) receptor, LDL receptor-related protein, and very low density lipoprotein (VLDL) receptor. Associates with high density lipoproteins (HDL) and the triacylglycerol-rich lipoproteins in the plasma and makes up about 10% of the protein of the VLDL and 2% of that of HDL. Appears to interfere directly with fatty acid uptake and is also the major plasma inhibitor of cholesteryl ester transfer protein (CETP). Binds free fatty acids and reduces their intracellular esterification. Modulates the interaction of APOE with beta-migrating VLDL and inhibits binding of beta-VLDL to the LDL receptor-related protein.

Research Area

Image Data



Western blot analysis of Apolipoprotein CI expression in Human plasma lysate.

Note

For research use only.