

Product Name: HIF-1 alpha (3M10) Rabbit Monoclonal Antibody
Catalog #: AMRe12017



Summary

Production Name	HIF-1 alpha (3M10) Rabbit Monoclonal Antibody
Description	Rabbit Monoclonal Antibody
Host	Rabbit
Application	WB,IHC-P,ICC/IF,FC,IP,IF-P
Reactivity	Human

Performance

Conjugation	Unconjugated
Modification	Unmodified
Isotype	IgG
Clonality	Monoclonal
Form	Liquid
Storage	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles. Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% New type
Buffer	preservative N and 50% glycerol. Store at +4°C short term. Store at -20°C long term. Avoid freeze / thaw cycle.
Purification	Affinity purification

Immunogen

Gene Name	HIF1A
Alternative Names	HIF1; MOP1; PASD8; bHLHe78; HIF-1alpha; HIF1-ALPHA; HIF1A
Gene ID	3091.0
SwissProt ID	Q16665.

Application

Dilution Ratio	WB 1:1000, IHC-P/IF-P 1:100, ICC/IF 1:200-1:500, FCM 1:500-1:2000, IP 1:50-1:200
Molecular Weight	93kDa

Product Name: HIF-1 alpha (3M10) Rabbit Monoclonal Antibody
Catalog #: AMRe12017



Background

Hypoxia-inducible factor-1 (HIF1) is a transcription factor found in mammalian cells cultured under reduced oxygen tension that plays an essential role in cellular and systemic homeostatic responses to hypoxia. HIF1 is a heterodimer composed of an alpha subunit and a beta subunit. The beta subunit has been identified as the aryl hydrocarbon receptor nuclear translocator (ARNT). This gene encodes the alpha subunit of HIF-1. Functions as a master transcriptional regulator of the adaptive response to hypoxia (PubMed:[11292861](http://www.uniprot.org/citations/11292861)), (PubMed:[11566883](http://www.uniprot.org/citations/11566883)), (PubMed:[15465032](http://www.uniprot.org/citations/15465032)), (PubMed:[16973622](http://www.uniprot.org/citations/16973622)), (PubMed:[17610843](http://www.uniprot.org/citations/17610843)), (PubMed:[18658046](http://www.uniprot.org/citations/18658046)), (PubMed:[20624928](http://www.uniprot.org/citations/20624928)), (PubMed:[22009797](http://www.uniprot.org/citations/22009797)), (PubMed:[9887100](http://www.uniprot.org/citations/9887100)), (PubMed:[30125331](http://www.uniprot.org/citations/30125331)). Under hypoxic conditions, activates the transcription of over 40 genes, including erythropoietin, glucose transporters, glycolytic enzymes, vascular endothelial growth factor, HILPDA, and other genes whose protein products increase oxygen delivery or facilitate metabolic adaptation to hypoxia (PubMed:[11292861](http://www.uniprot.org/citations/11292861)), (PubMed:[11566883](http://www.uniprot.org/citations/11566883)), (PubMed:[15465032](http://www.uniprot.org/citations/15465032)), (PubMed:[16973622](http://www.uniprot.org/citations/16973622)), (PubMed:[17610843](http://www.uniprot.org/citations/17610843)), (PubMed:[20624928](http://www.uniprot.org/citations/20624928)), (PubMed:[22009797](http://www.uniprot.org/citations/22009797)), (PubMed:[9887100](http://www.uniprot.org/citations/9887100)), (PubMed:[30125331](http://www.uniprot.org/citations/30125331)). Plays an essential role in embryonic vascularization, tumor angiogenesis and pathophysiology of ischemic disease (PubMed:[22009797](http://www.uniprot.org/citations/22009797)). Heterodimerizes with ARNT; heterodimer binds to core DNA sequence 5'-TACGTG-3' within the hypoxia response element (HRE) of target gene promoters (By similarity). Activation requires recruitment of transcriptional coactivators such as CREBBP and EP300 (PubMed:[9887100](http://www.uniprot.org/citations/9887100)), (PubMed:[16543236](http://www.uniprot.org/citations/16543236)). Activity is enhanced by interaction with NCOA1 and/or NCOA2 (PubMed:[10594042](http://www.uniprot.org/citations/10594042)). Interaction with redox regulatory protein APEX1 seems to activate CTAD and potentiates


Product Name: HIF-1 alpha (3M10) Rabbit Monoclonal Antibody
Catalog #: AMRe12017



activation by NCOA1 and CREBBP (PubMed: [10202154](http://www.uniprot.org/citations/10202154), PubMed: [10594042](http://www.uniprot.org/citations/10594042)). Involved in the axonal distribution and transport of mitochondria in neurons during hypoxia (PubMed: [19528298](http://www.uniprot.org/citations/19528298)).

Research Area

Image Data



Western blot analysis of HIF-1 alpha expression in Ramos cell lysate.

Note

For research use only.