# Product Name: Lin28B (12W5) Rabbit Monoclonal

**Antibody** 

Catalog #: AMRe13317



## **Summary**

Production Name Lin28B (12W5) Rabbit Monoclonal Antibody

**Description** Rabbit Monoclonal Antibody

**Host** Rabbit

**Application** WB,ICC/IF,FC,IP

Reactivity Human

#### **Performance**

ConjugationUnconjugatedModificationUnmodified

**Isotype** IgG

Clonality Monoclonal Form Liquid

**Storage** Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.

Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% New type

**Buffer** preservative N and 50% glycerol. Store at +4°C short term. Store at -20°C long term.

Avoid freeze / thaw cycle.

**Purification** Affinity purification

### **Immunogen**

Gene Name LIN28B

Alternative Names CSDD2; FLJ16517; LIN28B; Protein lin-28 homolog B;

 Gene ID
 389421.0

 SwissProt ID
 Q6ZN17.

# **Application**

**Dilution Ratio** WB 1:1000, ICC/IF 1:200, FCM 1:20, IP 1:20

Molecular Weight 27kDa

Web: https://www.enkilife.com E-mail: order@enkilife.com techsupport@enkilife.com Tel: 0086-27-87002838

Product Name: Lin28B (12W5) Rabbit Monoclonal

**Antibody** 

Catalog #: AMRe13317




## **Background**

It has recently been shown that upregulation of LIN28A and LIN28B in primary human tumors and human cancer cell lines is correlated with downregulation of let-7 miRNAs. LIN28 genes are reported to be involved in primordial germ cell development and germ cell malignancy. In addition, allelic variation in LIN28B is associated with regulating the timing of puberty in humans. Suppressor of microRNA (miRNA) biogenesis, including that of let-7 and possibly of miR107, miR-143 and miR-200c. Binds primary let-7 transcripts (pri-let-7), including pri-let-7g and pri-let-7a-1, and sequester them in the nucleolus, away from the microprocessor complex, hence preventing their processing into mature miRNA (PubMed: <a href="http://www.uniprot.org/citations/22118463" target=" blank">22118463</a>). Does not act on pri-miR21 (PubMed: <a href="http://www.uniprot.org/citations/22118463" target=" blank" > 22118463 </a >). The repression of let-7 expression is required for normal development and contributes to maintain the pluripotent state of embryonic stem cells by preventing let-7-mediated differentiation. When overexpressed, recruits ZCCHC11/TUT4 uridylyltransferase to pre-let-7 transcripts, leading to their terminal uridylation and degradation (PubMed:<a href="http://www.uniprot.org/citations/19703396" target=" blank">19703396</a>). This activity might not be relevant in vivo, as LIN28B-mediated inhibition of let-7 miRNA maturation appears to be ZCCHC11-independent (PubMed:<a href="http://www.uniprot.org/citations/22118463" target=" blank">22118463</a>). Interaction with target pre-miRNAs occurs via an 5'- GGAG-3' motif in the pre-miRNA terminal loop. Mediates MYC-induced let- 7 repression (By similarity). When overexpressed, isoform 1 stimulates growth of the breast adenocarcinoma cell line MCF-7. Isoform 2 has no effect on cell growth.

### **Research Area**

#### **Image Data**



Western blot analysis of Lin28B expression in K562 cell lysate.

#### Note

**Product Name: Lin28B (12W5) Rabbit Monoclonal** 

Antibody

Catalog #: AMRe13317



For research use only.

Web: https://www.enkilife.com E-mail: order@enkilife.com techsupport@enkilife.com Tel: 0086-27-87002838