

# Summary

Production Name	Serum Albumin (14W10) Rabbit Monoclonal Antibody
Description	Rabbit Monoclonal Antibody
Host	Rabbit
Application	WB
Reactivity	Human

# Performance

Conjugation	Unconjugated
Modification	Unmodified
lsotype	lgG
Clonality	Monoclonal
Form	Liquid
Storage	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.
	Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% New type
Buffer	preservative N and 50% glycerol. Store at +4°C short term. Store at -20°C long term.
	Avoid freeze / thaw cycle.
Purification	Affinity purification

#### Immunogen

Gene Name	ALB
Alternative Names	ALB; Albumin (32 AA) ;Albumin (AA 34) ;Albumin; BSA; Serum albumin; Bovine Serum
	Albumin
Gene ID	213.0
SwissProt ID	P02768.

# Application

Dilution Ratio	WB 1:1000-1:5000
Molecular Weight	69kDa




# Background

Albumin is a soluble, monomeric protein which comprises about one-half of the blood serum protein. Albumin functions primarily as a carrier protein for steroids, fatty acids, and thyroid hormones and plays a role in stabilizing extracellular fluid volume. Binds water, Ca(2+), Na(+), K(+), fatty acids, hormones, bilirubin and drugs (Probable). Its main function is the regulation of the colloidal osmotic pressure of blood (Probable). Major zinc transporter in plasma, typically binds about 80% of all plasma zinc (PubMed:<a href="http://www.uniprot.org/citations/19021548" target="\_blank">19021548</a>). Major calcium and magnesium transporter in plasma, binds approximately 45% of circulating calcium and magnesium in plasma (By similarity). Potentially has more than two calcium-binding sites and might additionally bind calcium in a non-specific manner (By similarity). The shared binding site between zinc and calcium at residue Asp-273 suggests a crosstalk between zinc and calcium transport in the blood (By similarity). The rank order of affinity is zinc > calcium > magnesium (By similarity). Binds to the bacterial siderophore enterobactin and inhibits enterobactin-mediated iron uptake of E.coli from ferric transferrin, and may thereby limit the utilization of iron and growth of enteric bacteria such as E.coli (PubMed:<a href="http://www.uniprot.org/citations/6234017" target="\_blank">6234017</a>). Does not prevent iron uptake by the bacterial siderophore aerobactin (PubMed:<a href="http://www.uniprot.org/citations/6234017" target="\_blank">6234017</a>). Does not prevent iron uptake by the bacterial siderophore aerobactin (PubMed:<a href="http://www.uniprot.org/citations/6234017" target="\_blank">6234017</a>). Does not prevent iron uptake by the bacterial siderophore aerobactin (PubMed:<a href="http://www.uniprot.org/citations/6234017" target="\_blank">6234017</a>). Does not prevent iron uptake by the bacterial siderophore aerobactin (PubMed:<a href="http://www.uniprot.org/citations/6234017" target="\_blank">6234017</a>).

### **Research Area**

### **Image Data**



Western blot analysis of Bovine Serum Albumin expression in (1) HeLa cell lysate; (2) NIH/3T3 cell lysate; (3) PC-12 cell lysate.

#### Note

For research use only.