Product Name: Tenascin C (7E16) Rabbit Monoclonal

Antibody

Catalog #: AMRe18788



Summary

Production Name Tenascin C (7E16) Rabbit Monoclonal Antibody

Description Rabbit Monoclonal Antibody

Host Rabbit

Application WB,IHC-P,IF-P **Reactivity** Human,Mouse,Rat

Performance

ConjugationUnconjugatedModificationUnmodified

Isotype IgG

Clonality Monoclonal Form Liquid

Storage Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.

Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% New type

Buffer preservative N and 50% glycerol. Store at +4°C short term. Store at -20°C long term.

Avoid freeze / thaw cycle.

Purification Affinity purification

Immunogen

Gene Name TNC

Tenascin; TN; Cytotactin; GMEM; GP 150-225; Glioma-associated-extracellular matrix

Alternative Names antigen; Hexabrachion; JI; Myotendinous antigen; Neuronectin; Tenascin C; TN-C; TNC;

HXB

 Gene ID
 3371.0

 SwissProt ID
 P24821.

Application

Dilution Ratio WB 1:1000-1:5000, IHC-P/IF-P 1:200-1:500

Molecular Weight 241kDa

Product Name: Tenascin C (7E16) Rabbit Monoclonal

Antibody

Catalog #: AMRe18788




Background

The tenascin family of extracellular matrix proteins includes Tenascin-C (also designated cytotactin or Tenascin), Tenascin-R (also designated restrictin, TN-R or janusin) and Tenascin-X. Tenascin proteins function as substrate-adhesion molecules (SAMs) and are involved in regulating numerous developmental processes, such as morphogenetic cell migration and organogenesis. The tenascin family proteins arise from various splicing events in the region of coding for FNIII repeats. Extracellular matrix protein implicated in guidance of migrating neurons as well as axons during development, synaptic plasticity as well as neuronal regeneration. Promotes neurite outgrowth from cortical neurons grown on a monolayer of astrocytes. Ligand for integrins alpha-8/beta-1, alpha-9/beta-1, alpha-V/beta-3 and alpha- V/beta-6. In tumors, stimulates angiogenesis by elongation, migration and sprouting of endothelial cells (PubMed:19884327).

Research Area

Image Data



Western blot analysis of Cytotactin C expression in (1)Human fetal brain lysate;(2)Human fetal kidney lysate.

Note

For research use only.

Web: https://www.enkilife.com E-mail: order@enkilife.com techsupport@enkilife.com Tel: 0086-27-87002838