

**Product Name: HER2 (Phospho Tyr1221/1222) Rabbit
Monoclonal Antibody
Catalog #: AMRe21108**



Summary

| | |
|------------------------|--|
| Production Name | HER2 (Phospho Tyr1221/1222) Rabbit Monoclonal Antibody |
| Description | Rabbit Monoclonal Antibody |
| Host | Rabbit |
| Application | WB,IF,IP,ELISA |
| Reactivity | Human |

Performance

| | |
|---------------------|--|
| Conjugation | Phospho |
| Modification | Phosphorylated |
| Isotype | IgG,Kappa |
| Clonality | Monoclonal |
| Form | Liquid |
| Storage | Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles. |
| Buffer | PBS, 50% glycerol, 0.05% Proclin 300, 0.05%BSA |
| Purification | Protein A |

Immunogen

| | |
|--------------------------|---|
| Gene Name | ERBB2 Receptor tyrosine-protein kinase erbB-2;Metastatic lymph node gene 19 protein;MLN |
| Alternative Names | 19;Proto-oncogene Neu;Proto-oncogene c-ErbB-2;Tyrosine kinase-type cell surface receptor HER2;p185erbB2;CD antigen CD340; |
| Gene ID | 2064.0 |
| SwissProt ID | P04626. |

Application

| | |
|-------------------------|---|
| Dilution Ratio | WB 1:2000-1:10000;IF 1:200-1:1000;ELISA 1:5000-1:20000;IP 1:50-1:200; |
| Molecular Weight | Calculated MW:138kD;Observed MW:185kD |

**Product Name: HER2 (Phospho Tyr1221/1222) Rabbit
Monoclonal Antibody
Catalog #: AMRe21108**




Background

Cell localization:Membrane.This gene encodes a member of the epidermal growth factor (EGF) receptor family of receptor tyrosine kinases. This protein has no ligand binding domain of its own and therefore cannot bind growth factors. However, it does bind tightly to other ligand-bound EGF receptor family members to form a heterodimer, stabilizing ligand binding and enhancing kinase-mediated activation of downstream signalling pathways, such as those involving mitogen-activated protein kinase and phosphatidylinositol-3 kinase. Allelic variations at amino acid positions 654 and 655 of isoform a (positions 624 and 625 of isoform b) have been reported, with the most common allele, Ile654/Ile655, shown here. Amplification and/or overexpression of this gene has been reported in numerous cancers, including breast and ovarian tumors. Alternative splicing results in several additional transcript variants, some encoding d

Research Area

Image Data



SK-BR-3 whole cell lysates were separated by 10% SDS-PAGE, and the membrane was blotted with primary antibody(1:1000).
The HRP-conjugated Goat anti-Rabbit IgG(H + L) antibody was used to detect the antibody.

Note

For research use only.