

**Product Name: BTK Rabbit Monoclonal Antibody**  
**Catalog #: AMRe21158**



## Summary

<b>Production Name</b>	BTK Rabbit Monoclonal Antibody
<b>Description</b>	Rabbit Monoclonal Antibody
<b>Host</b>	Rabbit
<b>Application</b>	WB,IHC,IF,IP,ELISA
<b>Reactivity</b>	Human

## Performance

<b>Conjugation</b>	Unconjugated
<b>Modification</b>	Unmodified
<b>Isotype</b>	IgG,Kappa
<b>Clonality</b>	Monoclonal
<b>Form</b>	Liquid
<b>Storage</b>	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.
<b>Buffer</b>	PBS, 50% glycerol, 0.05% Proclin 300, 0.05%BSA
<b>Purification</b>	Protein A

## Immunogen

<b>Gene Name</b>	BTK
<b>Alternative Names</b>	BTK;AGMX1;ATK;BPK;Tyrosine-protein kinase BTK;Agammaglobulinaemia tyrosine kinase;ATK;B-cell progenitor kinase;BPK;Bruton tyrosine kinase
<b>Gene ID</b>	695.0
<b>SwissProt ID</b>	Q06187.

## Application

<b>Dilution Ratio</b>	IHC 1:1000-1:4000;WB 1:2000-1:10000;IF 1:200-1:1000;ELISA 1:5000-1:20000;IP 1:50-1:200;
<b>Molecular Weight</b>	Calculated MW:67kD;Observed MW:67kD

## Background

**Product Name: BTK Rabbit Monoclonal Antibody**  
**Catalog #: AMRe21158**


---



Cell localization:Membrane,Cytoplasm.The protein encoded by this gene plays a crucial role in B-cell development. Mutations in this gene cause X-linked agammaglobulinemia type 1, which is an immunodeficiency characterized by the failure to produce mature B lymphocytes, and associated with a failure of Ig heavy chain rearrangement. Alternative splicing results in multiple transcript variants encoding different isoforms. [provided by RefSeq, Dec 2013],

## Research Area

## Image Data



RAW264.7 whole cell lysates were separated by 10% SDS-PAGE, and the membrane was blotted with primary antibody(1:1000). The HRP-conjugated Goat anti-Rabbit IgG(H + L) antibody was used to detect the antibody.

## Note

For research use only.