# Product Name: PKC $\alpha$ Rabbit Monoclonal Antibody

Catalog #: AMRe21177



## **Summary**

**Production Name** PKC α Rabbit Monoclonal Antibody

**Description** Rabbit Monoclonal Antibody

**Host** Rabbit

**Application** WB,IHC,IF,IP,ELISA **Reactivity** Human,Mouse,Rat

### **Performance**

ConjugationUnconjugatedModificationUnmodifiedIsotypeIgG,KappaClonalityMonoclonalFormLiquid

**Storage** Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.

Buffer PBS, 50% glycerol, 0.05% Proclin 300, 0.05%BSA

**Purification** Protein A

#### **Immunogen**

Gene Name PRKCA

Alternative Names PRKCA;PKCA;PRKACA;Protein kinase C alpha type;PKC-A;PKC-alpha

 Gene ID
 5578.0

 SwissProt ID
 P17252.

# **Application**

IHC 1:500-1:1000;WB 1:2000-1:10000;IF 1:200-1:1000;ELISA 1:5000-1:20000;IP 1:50-

**Dilution Ratio** 

1:200;

Molecular Weight Calculated MW:77kD;Observed MW:77kD

# **Background**

Cell localization:Membrane, Cytoplasm.Protein kinase C (PKC) is a family of serine- and threonine-specific protein kinases


# Product Name: PKC $\alpha$ Rabbit Monoclonal Antibody Catalog #: AMRe21177



that can be activated by calcium and the second messenger diacylglycerol. PKC family members phosphorylate a wide variety of protein targets and are known to be involved in diverse cellular signaling pathways. PKC family members also serve as major receptors for phorbol esters, a class of tumor promoters. Each member of the PKC family has a specific expression profile and is believed to play a distinct role in cells. The protein encoded by this gene is one of the PKC family members. This kinase has been reported to play roles in many different cellular processes, such as cell adhesion, cell transformation, cell cycle checkpoint, and cell volume control. Knockout studies in mice suggest that this kinase may be a fundamental regulator of cardiac contractility and Ca(2+) handling in myocytes. [provided by RefSeq, Jul 2

#### **Research Area**

#### **Image Data**



Hela-2 whole cell lysates were separated by 10% SDS-PAGE, and the membrane was blotted with primary antibody(1:1000).

The HRP-conjugated Goat anti-Rabbit IgG(H + L) antibody was used to detect the antibody.

#### Note

For research use only.