Product Name: ASK1 Rabbit Monoclonal Antibody

Catalog #: AMRe21269



Summary

Production Name ASK1 Rabbit Monoclonal Antibody

Description Rabbit Monoclonal Antibody

Host Rabbit

Application WB,IHC,IF,IP,ELISA **Reactivity** Human,Mouse

Performance

ConjugationUnconjugatedModificationUnmodifiedIsotypeIgG,KappaClonalityMonoclonalFormLiquid

Storage Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.

Buffer PBS, 50% glycerol, 0.05% Proclin 300, 0.05%BSA

Purification Protein A

Immunogen

Gene Name MAP3K5

MAP3K5;ASK1;MAPKKK5;MEKK5;Mitogen-activated protein kinase kinase kinase

Alternative Names 5; Apoptosis signal-regulating kinase 1; ASK-1; MAPK/ERK kinase kinase 5; MEK kinase

5;MEKK 5

Gene ID 4217.0 **SwissProt ID** Q99683.

Application

IHC 1:100-1:400;WB 1:1000-1:5000;IF 1:200-1:1000;ELISA 1:5000-1:20000;IP 1:50-

Dilution Ratio

1:200;

Molecular Weight Calculated MW:155kD;Observed MW:155kD

Product Name: ASK1 Rabbit Monoclonal Antibody

Catalog #: AMRe21269




Background

Cell localization:Cytoplasm.Mitogen-activated protein kinase (MAPK) signaling cascades include MAPK or extracellular signal-regulated kinase (ERK), MAPK kinase (MKK or MEK), and MAPK kinase kinase (MAPKK or MEKK). MAPKK kinase/MEKK phosphorylates and activates its downstream protein kinase, MAPK kinase/MEK, which in turn activates MAPK. The kinases of these signaling cascades are highly conserved, and homologs exist in yeast, Drosophila, and mammalian cells. MAPKKK5 contains 1,374 amino acids with all 11 kinase subdomains. Northern blot analysis shows that MAPKKK5 transcript is abundantly expressed in human heart and pancreas. The MAPKKK5 protein phosphorylates and activates MKK4 (aliases SERK1, MAPKK4) in vitro, and activates c-Jun N-terminal kinase (JNK)/stress-activated protein kinase (SAPK) during transient expression in COS and 293 cells; MAPKKK5 does not activate MAPK/ERK. [provided by Re

Research Area

Image Data



SiHa cell lysates were separated by 4-20% SDS-PAGE, and the membrane was blotted with primary antibody 1:1000. The HRP-conjugated Goat anti-Rabbit IgG(H + L) antibody was used to detect the antibody.

Note

For research use only.

Web: https://www.enkilife.com E-mail: order@enkilife.com techsupport@enkilife.com Tel: 0086-27-87002838