

Product Name: HDAC4 Rabbit Monoclonal Antibody
Catalog #: AMRe21270



Summary

Production Name	HDAC4 Rabbit Monoclonal Antibody
Description	Rabbit Monoclonal Antibody
Host	Rabbit
Application	WB,IHC,IF,IP,ELISA
Reactivity	Human,Mouse,Rat

Performance

Conjugation	Unconjugated
Modification	Unmodified
Isotype	IgG,Kappa
Clonality	Monoclonal
Form	Liquid
Storage	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.
Buffer	PBS, 50% glycerol, 0.05% Proclin 300, 0.05%BSA
Purification	Protein A

Immunogen

Gene Name	HDAC4
Alternative Names	HDAC4;KIAA0288;Histone deacetylase 4;HD4
Gene ID	9759.0
SwissProt ID	P56524.

Application

Dilution Ratio	IHC 1:50-1:200;WB 1:2000-1:10000;IF 1:200-1:1000;ELISA 1:5000-1:20000;IP 1:50-1:200;
Molecular Weight	Calculated MW:119kD;Observed MW:140kD

Background

Cell localization:Cytoplasm, Nucleus.Histones play a critical role in transcriptional regulation, cell cycle progression, and


Product Name: HDAC4 Rabbit Monoclonal Antibody
Catalog #: AMRe21270



developmental events. Histone acetylation/deacetylation alters chromosome structure and affects transcription factor access to DNA. The protein encoded by this gene belongs to class II of the histone deacetylase/acuc/apha family. It possesses histone deacetylase activity and represses transcription when tethered to a promoter. This protein does not bind DNA directly, but through transcription factors MEF2C and MEF2D. It seems to interact in a multiprotein complex with RbAp48 and HDAC3. [provided by RefSeq, Jul 2008],

Research Area

Image Data



HEK293 cell lysates were separated by 4-20% SDS-PAGE, and the membrane was blotted with primary antibody 1:1000. The HRP-conjugated Goat anti-Rabbit IgG(H + L) antibody was used to detect the antibody.

Note

For research use only.