

Product Name: Collagen IV Rabbit Monoclonal Antibody
Catalog #: AMRe21611



Summary

Production Name	Collagen IV Rabbit Monoclonal Antibody
Description	Rabbit Monoclonal Antibody
Host	Rabbit
Application	WB,IHC,IF,IP,ELISA
Reactivity	Human

Performance

Conjugation	Unconjugated
Modification	Unmodified
Isotype	IgG,Kappa
Clonality	Monoclonal
Form	Liquid
Storage	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.
Buffer	PBS, 50% glycerol, 0.05% Proclin 300, 0.05%BSA
Purification	Protein A

Immunogen

Gene Name	COL4A1 Arresten;BSVD;CO4A1_HUMAN;COL4A1;COL4A1 NC1 domain;COL4A2;COL4A3;COL4A4;COL4A5;collagen alpha-1;IV;chain;Collagen IV Alpha 1 Polypeptide;Collagen IV Alpha 2 Polypeptide;Collagen Of Basement Membrane Alpha 1 Chain;Collagen Of Basement Membrane Alpha 2 Chain;Collagen Type IV Alpha 1;collagen type IV alpha 1 chain;Collagen Type IV Alpha 2;Collagen Type IV Alpha 3;Collagen Type IV Alpha 4;Collagen Type IV Alpha 5;RatOR
Alternative Names	
Gene ID	1282.0
SwissProt ID	P02462.

Application

Dilution Ratio	IHC 1:200-1000;WB 1:1000-5000;IF 1:200-1000;ELISA 1:5000-20000;IP 1:50-200
-----------------------	--

Product Name: Collagen IV Rabbit Monoclonal Antibody
Catalog #: AMRe21611



Molecular Weight


Calculated MW:160kD;Observed MW:200kD

Background

Cell localization: Cytoplasmic. This gene encodes a type IV collagen alpha protein. Type IV collagen proteins are integral components of basement membranes. This gene shares a bidirectional promoter with a paralogous gene on the opposite strand. The protein consists of an amino-terminal 7S domain, a triple-helix forming collagenous domain, and a carboxy-terminal non-collagenous domain. It functions as part of a heterotrimer and interacts with other extracellular matrix components such as perlecan, proteoglycans, and laminins. In addition, proteolytic cleavage of the non-collagenous carboxy-terminal domain results in a biologically active fragment known as arresten, which has anti-angiogenic and tumor suppressor properties. Mutations in this gene cause porencephaly, cerebrovascular disease, and renal and muscular defects. Alternative splicing results in multiple transcript variants. [provided by RefSeq, Dec 2014],

Research Area


Image Data



Western blot analysis of lysates from HeLa

cells, using Collagen IV Rabbit mAb. The HRP-conjugated Goat anti-Rabbit IgG antibody was used to detect the antibody.

Product Name: Collagen IV Rabbit Monoclonal Antibody
Catalog #: AMRe21611



Immunohistochemical analysis of paraffin-embedded Human appendix tissue. 1, primary Antibody was diluted at 1:200(4°C,overnight). 2, EDTA pH 9.0 was used for antibody retrieval(>98°C,20min). 3,Secondary antibody was diluted at 1:200(room temperature, 30min).

Note

For research use only.