# Product Name: HLA DQB1/2 Rabbit Polyclonal Antibody Enkilife Catalog #: APRab00497

### **Summary**

**Production Name** HLA DQB1/2 Rabbit Polyclonal Antibody

**Description** Rabbit Polyclonal Antibody

**Host** Rabbit

**Application** WB,IHC-P,ELISA

**Reactivity** Human

### **Performance**

ConjugationUnconjugatedModificationUnmodified

**Isotype** IgG

ClonalityPolyclonalFormLiquid

Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw Storage

cycles.

**Buffer** Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide, pH 7.3.

**Purification** Affinity Purification

### **Immunogen**

Gene Name HLA-DQB1/HLA-DQB2

HLA-DQB1; HLA-DQB; HLA class II histocompatibility antigen; DQ beta 1 chain; MHC

class II antigen DQB1; HLA-DQB2; HLA-DXB; HLA class II histocompatibility antigen; DQ

Alternative Names

beta 2 chainHLA class II histocompatibility antigen; DX beta chain; MHC class II antigen

DQB2

**Gene ID** 3119/3120

**SwissProt ID** P01920/P05538.

# **Application**

**Dilution Ratio** WB: 1:500-1:1000 IHC: 1:50-1:100 ELISA: 1:10000

Molecular Weight Calculated MW: 30 kDa; Observed MW: 30 kDa

Web: https://www.enkilife.com E-mail: order@enkilife.com techsupport@enkilife.com Tel: 0086-27-87002838




# **Background**

Binds peptides derived from antigens that access the endocytic route of antigen presenting cells (APC) and presents them on the cell surface for recognition by the CD4 T-cells. The peptide binding cleft accommodates peptides of 10-30 residues. The peptides presented by MHC class II molecules are generated mostly by degradation of proteins that access the endocytic route, where they are processed by lysosomal proteases and other hydrolases.


### Research Area

Immunology

# **Image Data**




Western blot analysis of HLA DQB1/2 in PC-12 lysates using HLA DQB1/2 antibody.



Immunohistochemistry analysis of paraffin-embedded Human colon using HLA DQB1/2 antibody. High-pressure and temperature Sodium Citrate pH 6.0 was used for antigen retrieval.





Immunohistochemistry analysis of paraffin-embedded Human colon using HLA DQB1/2 antibody. High-pressure and temperature Sodium Citrate pH 6.0 was used for antigen retrieval.



Immunohistochemistry analysis of paraffin-embedded Human spleen using HLA DQB1/2 antibody. High-pressure and temperature Sodium Citrate pH 6.0 was used for antigen retrieval.

### Note

For research use only.