Product Name: Neuropilin 1 Rabbit Polyclonal Antibody Enkilife Catalog #: APRab00522

Summary

Production Name Neuropilin 1 Rabbit Polyclonal Antibody

Description Rabbit Polyclonal Antibody

Host Rabbit

Application WB,IHC-P,ELISA **Reactivity** Human,Mouse,Rat

Performance

ConjugationUnconjugatedModificationUnmodified

Isotype IgG

Clonality Polyclonal Form Liquid

Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw

cycles.

Buffer Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide, pH 7.3.

Purification Affinity Purification

Immunogen

Storage

Gene Name NRP1

NRP1; NRP; VEGF165R; Neuropilin-1; Vascular endothelial cell growth factor 165

receptor; CD antigen CD304

Gene ID 8829

SwissProt ID 014786.

Application

Dilution Ratio WB: 1:500-1:1000 IHC: 1:50-1:100 ELISA: 1:10000

Molecular Weight Calculated MW: 103 kDa; Observed MW: 100 kDa

Background




The membrane-bound isoform 1 is a receptor involved in the development of the cardiovascular system, in angiogenesis, in the formation of certain neuronal circuits and in organogenesis outside the nervous system. It mediates the chemorepulsant activity of semaphorins. It binds to semaphorin 3A, The PLGF-2 isoform of PGF, The VEGF-165 isoform of VEGF and VEGF-B. Coexpression with KDR results in increased VEGF-165 binding to KDR as well as increased chemotaxis. It may regulate VEGF-induced angiogenesis. The soluble isoform 2 binds VEGF-165 and appears to inhibit its binding to cells.


Research Area

Cardiovascular

Image Data




Western blot analysis of Neuropilin 1 in 293 lysates using Neuropilin 1 antibody.



Immunohistochemistry analysis of paraffin-embedded Human kidney using Neuropilin 1 antibody. High-pressure and temperature Sodium Citrate pH 6.0 was used for antigen retrieval.

Web: https://www.enkilife.com E-mail: order@enkilife.com techsupport@enkilife.com Tel: 0086-27-87002838





Immunohistochemistry analysis of paraffin-embedded Human kidney using Neuropilin 1 antibody. High-pressure and temperature Sodium Citrate pH 6.0 was used for antigen retrieval.

Note

For research use only.