

Summary

Cytochrome P450 2A6/7/13 Rabbit Polyclonal Antibody	
Rabbit Polyclonal Antibody	
Rabbit	
WB,ELISA	
Human	

Performance

Conjugation	Unconjugated	
Modification	Unmodified	
lsotype	lgG	
Clonality	Polyclonal	
Form	Liquid	
Storage	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.	
Buffer	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide, pH 7.3.	
Purification	Affinity Purification	

Immunogen

Gene Name	CYP2A6/CYP2A7/CYP2A13		
Alternative Names	CYP2A6; CYP2A3; Cytochrome P450 2A6; 1; 4-cineole 2-exo-monooxygenase; CYPIIA6;		
	Coumarin 7-hydroxylase; Cytochrome P450 IIA3; Cytochrome P450(I); CYP2A7;		
	Cytochrome P450 2A7; CYPIIA7; Cytochrome P450 IIA4; CYP2A13; Cytochrome P450		
	2A13; CYPIIA13		
Gene ID	1548/1549/1553		
SwissProt ID	P11509/P20853/Q16696.		

Application

Dilution Ratio	WB: 1:500-1:1000 ELISA: 1:10000
Molecular Weight	Calculated MW: 56 kDa; Observed MW: 56 kDa



Background

Exhibits a high coumarin 7-hydroxylase activity. Can act in the hydroxylation of the anti-cancer drugs cyclophosphamide and ifosphamide. Competent in the metabolic activation of aflatoxin B1. Constitutes the major nicotine C-oxidase. Acts as a 1,4-cineole 2-exo-monooxygenase. Possesses low phenacetin O-deethylation activity.

Research Area

Cardiovascular

Image Data



Western blot analysis of Cytochrome P450 2A6/7/13 in Jurkat lysates using CYP2A6/7/13 antibody.

Note

For research use only.