

**Product Name: ATG7 Rabbit Polyclonal Antibody**  
**Catalog #: APRab03731**



## Summary

<b>Production Name</b>	ATG7 Rabbit Polyclonal Antibody
<b>Description</b>	Rabbit Polyclonal Antibody
<b>Host</b>	Rabbit
<b>Application</b>	WB, ICC/IF
<b>Reactivity</b>	Human, Mouse, Rat

## Performance

<b>Conjugation</b>	Unconjugated
<b>Modification</b>	Unmodified
<b>Isotype</b>	IgG
<b>Clonality</b>	Polyclonal
<b>Form</b>	Liquid
<b>Storage</b>	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.
<b>Buffer</b>	Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.
<b>Purification</b>	Affinity Chromatography

## Immunogen

<b>Gene Name</b>	ATG7
<b>Alternative Names</b>	hAGP7; Ubiquitin-activating enzyme E1-like protein; APG7L
<b>Gene ID</b>	10533
<b>SwissProt ID</b>	O95352.

## Application

<b>Dilution Ratio</b>	WB: 1:500-1:1000 IF: 1:50-1:200
<b>Molecular Weight</b>	Calculated MW: 78 kDa; Observed MW: 78 kDa

## Background

The molecular machinery of autophagy was largely discovered in yeast and referred to as autophagy-related (Atg) genes.

**Product Name: ATG7 Rabbit Polyclonal Antibody**  
**Catalog #: AP Rab03731**




Formation of the autophagosome involves a ubiquitin-like conjugation system in which Atg12 is covalently bound to Atg5 and targeted to autophagosome vesicles. This conjugation reaction is mediated by the ubiquitin E1-like enzyme Atg7 and the E2-like enzyme Atg10.

## Research Area

Cell Biology

## Image Data



Western blot analysis of Atg7(Apg7) in (1) HepG2 lysates; (2) mouse spleen lysates; (3) rat kidney lysates using ATG7 antibody.

## Note

For research use only.