Product Name: ErbB-4 (phospho Tyr1284) Rabbit

Polyclonal Antibody Catalog #: APRab04628



Summary

Production Name ErbB-4 (phospho Tyr1284) Rabbit Polyclonal Antibody

Description Rabbit Polyclonal Antibody

Host Rabbit

Application WB,IHC-P,IF-P,IF-F,ICC/IF,ELISA

Reactivity Human, Mouse, Rat

Performance

Conjugation Unconjugated

Modification Phospho Antibody

Isotype IgG

Clonality Polyclonal Form Liquid

Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw Storage

cycles.

Buffer Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% New type preservative N.

Purification Affinity purification

Immunogen

Gene Name ERBB4

ERBB4; HER4; Receptor tyrosine-protein kinase erbB-4; Proto-oncogene-like protein c-Alternative Names

ErbB-4; Tyrosine kinase-type cell surface receptor HER4; p180erbB4

Gene ID 2066.0

Q15303.The antiserum was produced against synthesized peptide derived from human **SwissProt ID**

HER4 around the phosphorylation site of Tyr1284. AA range:1250-1299

Application

Dilution Ratio

WB 1:500-1:2000, IHC-P 1:100-1:300, IF-P/IF-F/ICC/IF 1:200-1:1000, ELISA 1:10000.Not

vot tostod in other application

yet tested in other applications.

Web: https://www.enkilife.com E-mail: order@enkilife.com techsupport@enkilife.com Tel: 0086-27-87002838

Product Name: ErbB-4 (phospho Tyr1284) Rabbit

Polyclonal Antibody Catalog #: APRab04628



Molecular Weight

180kDa


Background

This gene is a member of the Tyr protein kinase family and the epidermal growth factor receptor subfamily. It encodes a single-pass type I membrane protein with multiple cysteine rich domains, a transmembrane domain, a tyrosine kinase domain, a phosphotidylinositol-3 kinase binding site and a PDZ domain binding motif. The protein binds to and is activated by neuregulins and other factors and induces a variety of cellular responses including mitogenesis and differentiation. Multiple proteolytic events allow for the release of a cytoplasmic fragment and an extracellular fragment. Mutations in this gene have been associated with cancer. Alternatively spliced variants which encode different protein isoforms have been described; however, not all variants have been fully characterized. [provided by RefSeq, Jul 2008],catalytic activity:ATP + a [protein]-L-tyrosine = ADP + a [protein]-L-tyrosine phosphate.,domain:The WW-binding motifs mediate interaction with WWOX., function: Specifically binds and is activated by neuregulins, NRG-2, NRG-3, heparinbinding EGF-like growth factor, betacellulin and NTAK. Interaction with these factors induces cell differentiation. Not activated by EGF, TGF-A, and amphiregulin., PTM: Isoform JM-A is processed but not isoform JM-B. So, they respectively represent cleavable and non-cleavable forms of the receptor, PTM: Ligand-binding increases phosphorylation on tyrosine residues, similarity: Belongs to the protein kinase superfamily. Tyr protein kinase family. EGF receptor subfamily, similarity: Contains 1 protein kinase domain, subunit: Homodimer or heterodimer with each of the other ERBB receptors (Potential). Interacts with PDZ domains of DLG2, DLG3, DLG4 and the syntrophin SNTB2. Interacts with CBFA2T3, MUC1 and WWOX.,tissue specificity:Expressed at highest levels in brain, heart, kidney, in addition to skeletal muscle, parathyroid, cerebellum, pituitary, spleen, testis and breast. Lower levels in thymus, lung, salivary gland, and pancreas. Isoform JM-A and isoform JM-B are expressed in cerebellum, but only the isoform JM-B is expressed in the heart.,

Research Area

ErbB_HER;Calcium;Endocytosis;

Image Data




Web: https://www.enkilife.com E-mail: order@enkilife.com techsupport@enkilife.com Tel: 0086-27-87002838

Product Name: ErbB-4 (phospho Tyr1284) Rabbit

Polyclonal Antibody Catalog #: APRab04628




Immunofluorescence analysis of HeLa cells treated with EGF 200nM 5 ', using HER4 (Phospho-Tyr1284) Antibody. The picture on the right is blocked with the phospho peptide.



Immunohistochemistry analysis of paraffin-embedded human breast carcinoma, using HER4 (Phospho-Tyr1284) Antibody.

The picture on the right is blocked with the phospho peptide.



Western blot analysis of lysates from HUVEC cells treated with EGF 200ng/ml 30 ', using HER4 (Phospho-Tyr1284)

Antibody. The lane on the right is blocked with the phospho peptide.

Note

For research use only.