Polyclonal Antibody Catalog #: APRab04695



### **Summary**

**Production Name** FoxO1 (phospho Ser319) Rabbit Polyclonal Antibody

**Description** Rabbit Polyclonal Antibody

**Host** Rabbit

**Application** WB,IHC-P,IF-P,IF-F,ICC/IF,ELISA

**Reactivity** Human, Mouse, Rat

#### **Performance**

**Conjugation** Unconjugated

**Modification** Phospho Antibody

**Isotype** IgG

Clonality Polyclonal Form Liquid

Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw Storage

cycles.

**Buffer** Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% New type preservative N.

**Purification** Affinity purification

#### **Immunogen**

Gene Name FOXO1

FOXO1; FKHR; FOXO1A; Forkhead box protein O1; Forkhead box protein O1A; Forkhead Alternative Names

in rhabdomyosarcoma

**Gene ID** 2308.0

Q12778.The antiserum was produced against synthesized peptide derived from human **SwissProt ID** 

FKHR around the phosphorylation site of Ser319. AA range:286-335

### **Application**

**Dilution Ratio** WB 1:500-1:2000, IHC-P 1:100-1:300, ELISA 1:10000, IF-P/IF-F/ICC/IF 1:50-200

Molecular Weight 82kDa

Polyclonal Antibody Catalog #: APRab04695




### **Background**

This gene belongs to the forkhead family of transcription factors which are characterized by a distinct forkhead domain. The specific function of this gene has not yet been determined; however, it may play a role in myogenic growth and differentiation. Translocation of this gene with PAX3 has been associated with alveolar rhabdomyosarcoma. [provided by RefSeq, Jul 2008], disease: Chromosomal aberrations involving FOXO1 are a cause of rhabdomyosarcoma 2 (RMS2) [MIM:268220]; also known as alveolar rhabdomyosarcoma. Translocation (2;13)(q35;q14) with PAX3; translocation t(1;13) (p36;q14) with PAX7. The resulting protein is a transcriptional activator, function: Transcription factor, PTM: Phosphorylated by AKT1; insulin-induced (By similarity). IGF1 rapidly induces phosphorylation of Ser-256, Thr-24, and Ser-319. Phosphorylation of Ser-256 decreases DNA-binding activity and promotes the phosphorylation of Thr-24, and Ser-319, permitting phosphorylation of Ser-322 and Ser-325, probably by CK1, leading to nuclear exclusion and loss of function. Phosphorylation of Ser-329 is independent of IGF1 and leads to reduced function. Phosphorylated upon DNA damage, probably by ATM or ATR, similarity: Contains 1 fork-head DNA-binding domain, subcellular location: Shuttles between cytoplasm and nucleus, subunit: Interacts with LRPPRC, tissue specificity: Ubiquitous,

#### **Research Area**

Insulin Receptor; B Cell Receptor; Protein Acetylation

### **Image Data**




Enzyme-Linked Immunosorbent Assay (Phospho-ELISA) for Immunogen Phosphopeptide (Phospho-left) and Non-Phosphopeptide (Phospho-right) , using FKHR (Phospho-Ser319) Antibody

Web: https://www.enkilife.com E-mail: order@enkilife.com techsupport@enkilife.com Tel: 0086-27-87002838


Polyclonal Antibody Catalog #: APRab04695






Immunohistochemistry analysis of paraffin-embedded human breast carcinoma, using FKHR (Phospho-Ser319) Antibody.

The picture on the right is blocked with the phospho peptide.




Western blot analysis of lysates from HeLa cells treated with EGF, using FKHR (Phospho-Ser319) Antibody. The lane on the right is blocked with the phospho peptide.



Western Blot analysis of various cells using Phospho-FoxO1 (S319) Polyclonal Antibody

Polyclonal Antibody Catalog #: APRab04695





Western Blot analysis of 293 cells using Phospho-FoxO1 (S319) Polyclonal Antibody

#### Note

For research use only.