



Catalog #: APRab05052

## Summary

Myf-5 (phospho Ser49) Rabbit Polyclonal Antibody **Production Name** 

Description Rabbit Polyclonal Antibody

Host Rabbit

**Application** WB,IHC-P,IF-P,IF-F,ICC/IF,ELISA

Reactivity Human, Mouse

#### **Performance**

Conjugation Unconjugated

Phospho Antibody Modification

Isotype IgG

**Clonality** Polyclonal **Form** Liquid

Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw Storage

cycles.

**Buffer** Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% New type preservative N.

**Purification** Affinity purification

### **Immunogen**

Gene Name MYF5

MYF5; BHLHC2; Myogenic factor 5; Myf-5; Class C basic helix-loop-helix protein 2; **Alternative Names** 

bHLHc2

Gene ID 4617.0

P13349.The antiserum was produced against synthesized peptide derived from human SwissProt ID

Myogenic Factor 5 around the phosphorylation site of Ser49. AA range:21-70

# **Application**

**Dilution Ratio** WB 1:500-1:2000, IHC-P 1:100-1:300, ELISA 1:10000, IF-P/IF-F/ICC/IF 1:50-200

**Molecular Weight** 38kDa

Web: https://www.enkilife.com E-mail: order@enkilife.com techsupport@enkilife.com Tel: 0086-27-87002838



Catalog #: APRab05052




## **Background**


function:Involved in muscle differentiation (myogenic factor). Induces fibroblasts to differentiate into myoblasts. Probable sequence specific DNA-binding protein., similarity: Contains 1 basic helix-loop-helix (bHLH) domain., subunit: Efficient DNA binding requires dimerization with another bHLH protein., function: Involved in muscle differentiation (myogenic factor). Induces fibroblasts to differentiate into myoblasts. Probable sequence specific DNA-binding protein., similarity: Contains 1 basic helix-loop-helix (bHLH) domain., subunit: Efficient DNA binding requires dimerization with another bHLH protein.,

#### Research Area

# **Image Data**



Western blot analysis of lysates from HeLa cells, using Myogenic Factor 5 (Phospho-Ser49) Antibody. The lane on the right is blocked with the phospho peptide.



Immunohistochemical analysis of paraffin-embedded Human breast cancer. Antibody was diluted at 1:100 (4°,overnight) . High-pressure and temperature Tris-EDTA,pH8.0 was used for antigen retrieval. Negetive contrl (right) obtaned from antibody was pre-absorbed by immunogen peptide.

Web: https://www.enkilife.com E-mail: order@enkilife.com techsupport@enkilife.com Tel: 0086-27-87002838





Catalog #: APRab05052



### Note

For research use only.

Tel: 0086-27-87002838