Product Name: ACSS1 (Acetyl-K642) Rabbit Polyclonal



Catalog #: APRab06169



Summary

Production Name ACSS1 (Acetyl-K642) Rabbit Polyclonal Antibody

Description Rabbit Polyclonal Antibody

Host Rabbit
Application WB,ELISA

Reactivity Human, Mouse, Rat

Performance

ConjugationUnconjugatedModificationAcetyl Antibody

Isotype IgG

Clonality Polyclonal Form Liquid

Storage Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.

Buffer Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% New type preservative N.

Purification Affinity purification

Immunogen

Gene Name ACSS1 ACAS2L KIAA1846

Acetyl-coenzyme A synthetase 2-like, mitochondrial (EC 6.2.1.1) (Acetate--CoA ligase 2)

Alternative Names (Acetyl-CoA synthetase 2) (AceCS2) (Acyl-CoA synthetase short-chain family member

1)

Gene ID 84532.0

SwissProt ID Q9NUB1. Synthesized Acetyl peptide derived from human ACSS1. at AA range: K642

Application

Dilution Ratio WB 1:500-1:2000, ELISA 1:20000.Not yet tested in other applications.

Molecular Weight

Web: https://www.enkilife.com E-mail: order@enkilife.com techsupport@enkilife.com Tel: 0086-27-87002838

Product Name: ACSS1 (Acetyl-K642) Rabbit Polyclonal

Antibody

Catalog #: APRab06169




Background

This gene encodes a mitochondrial acetyl-CoA synthetase enzyme. A similar protein in mice plays an important role in the tricarboxylic acid cycle by catalyzing the conversion of acetate to acetyl CoA. Alternatively spliced transcript variants encoding multiple isoforms have been observed for this gene. [provided by RefSeq, Nov 2011],catalytic activity:ATP + acetate + CoA = AMP + diphosphate + acetyl-CoA.,function:Converts acetate to acetyl-CoA so that it can be used for oxidation through the tricarboxylic cycle to produce ATP and CO(2).,sequence caution:Sequencing errors.,similarity:Belongs to the ATP-dependent AMP-binding enzyme family.,

Research Area

Glycolysis / Gluconeogenesis;Pyruvate metabolism;Propanoate metabolism;

Image Data



Western Blot analysis of 1,293T 2,hela cells using primary antibody diluted at 1:1000 (4°C overnight) . Secondary antibody: Goat Anti-rabbit IgG IRDye 800 (diluted at 1:5000, 25°C, 1 hour)

Note

For research use only.

Web: https://www.enkilife.com E-mail: order@enkilife.com techsupport@enkilife.com Tel: 0086-27-87002838