

Product Name: Histone H2B (Acetyl Lys126) Rabbit Polyclonal Antibody
Catalog #: APRab06195



Summary

Production Name	Histone H2B (Acetyl Lys126) Rabbit Polyclonal Antibody
Description	Rabbit Polyclonal Antibody
Host	Rabbit
Application	WB,ELISA
Reactivity	Human,Mouse,Rat

Performance

Conjugation	Unconjugated
Modification	Acetyl Antibody
Isotype	IgG
Clonality	Polyclonal
Form	Liquid
Storage	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.
Buffer	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% New type preservative N.
Purification	Affinity purification

Immunogen

Gene Name	HIST1H2BC HIST1H2BA; TSH2B; Histone H2B type 1-A; Histone H2B, testis; Testis-specific histone H2B; HIST1H2BB; H2BFF; Histone H2B type 1-B; Histone H2B.1; Histone H2B.f; H2B/f; HIST1H2BC; H2BFL; HIST1H2BE; H2BFH; HIST1H2BF;H2BFG; HIST1H2BG; H2BFA; HIST1H2BI; H2BFK; Histone H2B type 1-C/E/F/G/I; Histone H2B.1 A; Histone H2B.a; H2B/a; Histone H2B.g; H2B/g; Histone H2B.h; H2B/h; Histone H2B.k; H2B/k; Histone H2B.l; H2B/l;H2BK126AC
Alternative Names	
Gene ID	255626.0
SwissProt ID	Q96A08.The antiserum was produced against synthesized Acetyl-peptide derived from human H2B around the Acetylation site of Lys126. AA range:78-127

Application

Product Name: Histone H2B (Acetyl Lys126) Rabbit Polyclonal Antibody
Catalog #: APRab06195



Dilution Ratio	WB 1:500-1:2000, ELISA 1:20000. Not yet tested in other applications.
Molecular Weight	15kDa


Background

Histones are basic nuclear proteins that are responsible for the nucleosome structure of the chromosomal fiber in eukaryotes. Nucleosomes consist of approximately 146 bp of DNA wrapped around a histone octamer composed of pairs of each of the four core histones (H2A, H2B, H3, and H4). The chromatin fiber is further compacted through the interaction of a linker histone, H1, with the DNA between the nucleosomes to form higher order chromatin structures. This gene is intronless and encodes a replication-dependent histone that is a testis/sperm-specific member of the histone H2B family. Transcripts from this gene contain a palindromic termination element. [provided by RefSeq, Aug 2015], function: Core component of nucleosome. Nucleosomes wrap and compact DNA into chromatin, limiting DNA accessibility to the cellular machineries which require DNA as a template. Histones thereby play a central role in transcription regulation, DNA repair, DNA replication and chromosomal stability. DNA accessibility is regulated via a complex set of post-translational modifications of histones, also called histone code, and nucleosome remodeling., PTM: Monoubiquitination of Lys-122 by the RNF20/40 complex gives a specific tag for epigenetic transcriptional activation and is also prerequisite for histone H3 'Lys-4' and 'Lys-79' methylation. It also functions cooperatively with the FACT dimer to stimulate elongation by RNA polymerase II., similarity: Belongs to the histone H2B family., subunit: The nucleosome is a histone octamer containing two molecules each of H2A, H2B, H3 and H4 assembled in one H3-H4 heterotetramer and two H2A-H2B heterodimers. The octamer wraps approximately 147 bp of DNA., tissue specificity: Transcribed exclusively in testis, and the corresponding protein is also present in mature sperm.,

Research Area

Systemic lupus erythematosus;

Image Data




**Product Name: Histone H2B (Acetyl Lys126) Rabbit
Polyclonal Antibody
Catalog #: APRab06195**



Western blot analysis of lysate from A549 cells, using H2B (Acetyl-Lys126) Antibody.



Western Blot analysis of A549 cells using Acetyl-Histone H2B (K126) Polyclonal Antibody.. Secondary antibody was diluted at 1:20000



Western Blot analysis of A549 cells using Acetyl-Histone H2B (K126) Polyclonal Antibody. Secondary antibody was diluted at 1:20000

Note

For research use only.