Product Name: TAL1/2 (Acetyl Lys221/Acetyl Lys222/Acetyl Lys36/Acetyl Lys37) Rabbit Polyclonal



Antibody

Catalog #: APRab06260

Summary

Production Name TAL1/2 (Acetyl Lys221/Acetyl Lys26/Acetyl Lys36/Acetyl Lys37) Rabbit Polyclonal

Antibody

Description Rabbit Polyclonal Antibody

Host Rabbit

Application WB,ELISA

Reactivity Human, Mouse, Rat

Performance

ConjugationUnconjugatedModificationAcetyl Antibody

Isotype IgG

Clonality Polyclonal Form Liquid

Storage Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.

Buffer Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% New type preservative N.

Purification Affinity purification

Immunogen

Alternative Names

Gene Name TAL1/TAL2

TAL1; BHLHA17; SCL; TCL5; T-cell acute lymphocytic leukemia protein 1; TAL-1; Class A

basic helix-loop-helix protein 17; bHLHa17; Stem cell protein; T-cell

leukemia/lymphoma protein 5; TAL2; BHLHA19; T-cell acute lymphocytic leukemia

protein 2; TAL-2; Class A basic helix-loop-helix protein 19; bHLHa19

Gene ID 6886.0

P17542. Synthesized acetyl-peptide derived from human TAL1/2 around the SwissProt ID

acetylation site of K221.

Application

Web: https://www.enkilife.com E-mail: order@enkilife.com techsupport@enkilife.com Tel: 0086-27-87002838

Product Name: TAL1/2 (Acetyl Lys221/Acetyl Lys222/Acetyl Lys36/Acetyl Lys37) Rabbit Polyclonal Antibody



Catalog #: APRab06260

Dilution Ratio WB 1:500-1:2000, ELISA 1:10000.Not yet tested in other applications.

Molecular Weight 45kDa

Background

alternative products: The splicing pattern is cell-lineage dependent, disease: A chromosomal aberration involving TAL1 may be a cause of some T-cell acute lymphoblastic leukemias (T-ALL). Translocation t(1;14)(p32;q11) with T-cell receptor alpha chain (TCRA) genes., domain: The helix-loop-helix domain is necessary and sufficient for the interaction with DRG1, function: Implicated in the genesis of hemopoietic malignancies. It may play an important role in hemopoietic differentiation. Serves as a positive regulator of erythroid differentiation, PTM: Phosphorylated on serine residues. Phosphorylation of Ser-122 is strongly stimulated by hypoxia., PTM: Ubiquitinated; subsequent to hypoxia-dependent phosphorylation of Ser-122, ubiquitination targets the protein for rapid degradation via the ubiquitin system. This process may be characteristic for microvascular endothelial cells, since it could not be observed in large vessel endothelial cells,,similarity:Contains 1 basic helix-loop-helix (bHLH) domain,,subunit:Efficient DNA binding requires dimerization with another bHLH protein. Forms heterodimers with TCF3. Binds to the LIM domain containing protein LMO2 and to DRG1. Can assemble in a complex with LDB1 and LMO2. Component of a TAL-1 complex composed at least of CBFA2T3, LDB1, TAL1 and TCF3.,tissue specificity:Leukemic stem cell.,alternative products:The splicing pattern is cell-lineage dependent,disease:A chromosomal aberration involving TAL1 may be a cause of some T-cell acute lymphoblastic leukemias (T-ALL). Translocation t(1;14)(p32;q11) with T-cell receptor alpha chain (TCRA) genes.,domain:The helix-loop-helix domain is necessary and sufficient for the interaction with DRG1, function: Implicated in the genesis of hemopoietic malignancies. It may play an important role in hemopoietic differentiation. Serves as a positive regulator of erythroid differentiation.,PTM:Phosphorylated on serine residues. Phosphorylation of Ser-122 is strongly stimulated by hypoxia., PTM: Ubiquitinated; subsequent to hypoxia-dependent phosphorylation of Ser-122, ubiquitination targets the protein for rapid degradation via the ubiquitin system. This process may be characteristic for microvascular endothelial cells, since it could not be observed in large vessel endothelial cells, similarity: Contains 1 basic helix-loop-helix (bHLH) domain., subunit: Efficient DNA binding requires dimerization with another bHLH protein. Forms heterodimers with TCF3. Binds to the LIM domain containing protein LMO2 and to DRG1. Can assemble in a complex with LDB1 and LMO2. Component of a TAL-1 complex composed at least of CBFA2T3, LDB1, TAL1 and TCF3., tissue specificity: Leukemic stem cell.,

Research Area


Image Data

Web: https://www.enkilife.com E-mail: order@enkilife.com techsupport@enkilife.com Tel: 0086-27-87002838

C EnkiLife

Product Name: TAL1/2 (Acetyl Lys221/Acetyl Lys222/Acetyl Lys36/Acetyl Lys37) Rabbit Polyclonal Antibody

Catalog #: APRab06260



Western blot analysis of MOUSE-HEART MOUSE-BRAIN using Acetyl-TAL1/2 (K221/K222/K36/K37) antibody. Antibody was diluted at 1:500. Secondary antibody was diluted at 1:20000

Note

For research use only.

Web: https://www.enkilife.com E-mail: order@enkilife.com techsupport@enkilife.com Tel: 0086-27-87002838