Product Name: Tubulin α (Acetyl Lys352) Rabbit

Polyclonal Antibody Catalog #: APRab06264



Summary

Tubulin α (Acetyl Lys352) Rabbit Polyclonal Antibody **Production Name**

Description Rabbit Polyclonal Antibody

Host Rabbit **Application** WB.ELISA

Reactivity Human, Mouse, Rat

Performance

Unconjugated Conjugation Modification Acetyl Antibody

Isotype IgG

Clonality Polyclonal **Form** Liquid

Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw Storage

cycles.

Buffer Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% New type preservative N.

Purification Affinity purification

Immunogen

Alternative Names

Gene Name TUBA1A/TUBA1B/TUBA1C/TUBA3C/TUBA4A/TUBA8

> TUBA1A; TUBA3; Tubulin alpha-1A chain; Alpha-tubulin 3; Tubulin B-alpha-1; Tubulin alpha-3 chain; TUBA1B; Tubulin alpha-1B chain; Alpha-tubulin ubiquitous; Tubulin K-

> alpha-1; Tubulin alpha-ubiquitous chain; TUBA1C; TUBA6;Tubulin alpha-6 chain;

TUBA3C; TUBA3D; Tubulin alpha-3C/D chain; Alpha-tubulin 2; Alpha-tubulin 3C/D; Tubulin alpha-2 chain; TUBA4A; TUBA1; Tubulin alpha-4A chain; Alpha-tubulin 1;

Testis-specific alpha-tubulin;

Gene ID 7846.0

Q71U36. The antiserum was produced against synthesized Acetyl-peptide derived from SwissProt ID

human TUBA1B around the Acetylation site of Lys352. AA range:311-360

Application

Product Name: Tubulin α (Acetyl Lys352) Rabbit

Polyclonal Antibody Catalog #: APRab06264



Dilution Ratio WB 1:500-1:2000, ELISA 1:20000.Not yet tested in other applications.

Molecular Weight 50kDa

Background

Microtubules of the eukaryotic cytoskeleton perform essential and diverse functions and are composed of a heterodimer of alpha and beta tubulins. The genes encoding these microtubule constituents belong to the tubulin superfamily, which is composed of six distinct families. Genes from the alpha, beta and gamma tubulin families are found in all eukaryotes. The alpha and beta tubulins represent the major components of microtubules, while gamma tubulin plays a critical role in the nucleation of microtubule assembly. There are multiple alpha and beta tubulin genes, which are highly conserved among species. This gene encodes alpha tubulin and is highly similar to the mouse and rat Tuba1 genes. Northern blotting studies have shown that the gene expression is predominantly found in morphologically differentiated neurologic cells. This gene is one of three alpha-tubulin genes in a cluster on chromosome 12q.disease:Defects in TUBA1A are the cause of lissencephaly type 3 (LIS3) [MIM:611603]. LIS is characterized by a smooth brain surface due to the absence (agyria) or reduction (pachygyria) of surface convolutions. It is often associated with psychomotor retardation and seizures. LIS3 features include agyria or pachygyria or laminar heterotopia, severe mental retardation, motor delay, variable presence of seizures, and abnormalities of corpus callosum, hippocampus, cerebellar vermis and brainstem.,function:Tubulin is the major constituent of microtubules. It binds two moles of GTP, one at an exchangeable site on the beta chain and one at a non-exchangeable site on the alpha-chain., PTM: Undergoes a tyrosination/detyrosination cycle, the cyclic removal and readdition of a C-terminal tyrosine residue by the enzymes tubulin tyrosine carboxypeptidase (TTCP) and tubulin tyrosine ligase (TTL), respectively., similarity: Belongs to the tubulin family., subunit: Dimer of alpha and beta chains., tissue specificity:Expressed at a high level in fetal brain.,

Research Area

Gap junction; Pathogenic Escherichia coli infection;


Image Data

Web: https://www.enkilife.com E-mail: order@enkilife.com techsupport@enkilife.com Tel: 0086-27-87002838


Product Name: Tubulin α (Acetyl Lys352) Rabbit

Polyclonal Antibody Catalog #: APRab06264






Western blot analysis of lysate from K562 cells, using TUBA1B (Acetyl-Lys352) Antibody.



Western Blot analysis of K562, A549, Jurkat cells using Acetyl-Tubulin α (K352) Polyclonal Antibody. Secondary antibody was diluted at 1:20000



Western Blot analysis of K562 cells using Acetyl-Tubulin α (K352) Polyclonal Antibody. Secondary antibody was diluted at 1:20000

Note

For research use only.