
Summary

Production Name	ADAMTS-1 Rabbit Polyclonal Antibody
Description	Rabbit Polyclonal Antibody
Host	Rabbit
Application	WB,ELISA
Reactivity	Human,Rat,Mouse

Performance

Conjugation	Unconjugated
Modification	Unmodified
Isotype	IgG
Clonality	Polyclonal
Form	Liquid
Storage	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.
Buffer	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% New type preservative N.
Purification	Affinity purification

Immunogen

Gene Name	ADAMTS1
Alternative Names	ADAMTS1; KIAA1346; METH1; A disintegrin and metalloproteinase with thrombospondin motifs 1; ADAM-TS 1; ADAM-TS1; ADAMTS-1; METH-1
Gene ID	9510.0
SwissProt ID	Q9UHI8.Synthesized peptide derived from ADAMTS-1 . at AA range: 160-240

Application

Dilution Ratio	WB 1:500-1:2000, ELISA 1:5000.Not yet tested in other applications.
Molecular Weight	105kDa

Background


Product Name: ADAMTS-1 Rabbit Polyclonal Antibody
Catalog #: APRab06595



This gene encodes a member of the ADAMTS (a disintegrin and metalloproteinase with thrombospondin motif) protein family. Members of the family share several distinct protein modules, including a propeptide region, a metalloproteinase domain, a disintegrin-like domain, and a thrombospondin type 1 (TS) motif. Individual members of this family differ in the number of C-terminal TS motifs, and some have unique C-terminal domains. The protein encoded by this gene contains two disintegrin loops and three C-terminal TS motifs and has anti-angiogenic activity. The expression of this gene may be associated with various inflammatory processes as well as development of cancer cachexia. This gene is likely to be necessary for normal growth, fertility, and organ morphology and function. [provided by RefSeq, Jul 2008],catalytic activity:Cleaves aggrecan at the 1938-Glu-|-Leu-1939 site, within the chondroitin sulfate attachment domain.,cofactor:Binds 1 zinc ion per subunit.,domain:The conserved cysteine present in the cysteine-switch motif binds the catalytic zinc ion, thus inhibiting the enzyme. The dissociation of the cysteine from the zinc ion upon the activation-peptide release activates the enzyme.,domain:The spacer domain and the TSP type-1 domains are important for a tight interaction with the extracellular matrix.,function:Cleaves aggrecan, a cartilage proteoglycan, and may be involved in its turnover (By similarity). Has angiogenic inhibitor activity. Active metalloprotease, which may be associated with various inflammatory processes as well as development of cancer cachexia. May play a critical role in follicular rupture.,PTM:The precursor is cleaved by a furin endopeptidase.,similarity:Contains 1 disintegrin domain.,similarity:Contains 1 peptidase M12B domain.,similarity:Contains 3 TSP type-1 domains.,


Research Area

Image Data



Western Blot analysis of various cells using ADAMTS-1 Polyclonal Antibody

Product Name: ADAMTS-1 Rabbit Polyclonal Antibody
Catalog #: APRab06595



Western Blot analysis of HUVEC cells using ADAMTS-1 Polyclonal Antibody

Note

For research use only.