

**Product Name: ADAMTS-18 Rabbit Polyclonal Antibody**  
**Catalog #: APRab06599**



## Summary

<b>Production Name</b>	ADAMTS-18 Rabbit Polyclonal Antibody
<b>Description</b>	Rabbit Polyclonal Antibody
<b>Host</b>	Rabbit
<b>Application</b>	WB,IHC-P,IF-P,IF-F,ICC/IF,ELISA
<b>Reactivity</b>	Human,Mouse

## Performance

<b>Conjugation</b>	Unconjugated
<b>Modification</b>	Unmodified
<b>Isotype</b>	IgG
<b>Clonality</b>	Polyclonal
<b>Form</b>	Liquid
<b>Storage</b>	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.
<b>Buffer</b>	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% New type preservative N.
<b>Purification</b>	Affinity purification

## Immunogen

<b>Gene Name</b>	ADAMTS18
<b>Alternative Names</b>	ADAMTS18; ADAMTS21; A disintegrin and metalloproteinase with thrombospondin motifs 18; ADAM-TS 18; ADAM-TS18; ADAMTS-18
<b>Gene ID</b>	170692.0
<b>SwissProt ID</b>	Q8TE60.Synthesized peptide derived from ADAMTS-18 . at AA range: 1030-1110

## Application

<b>Dilution Ratio</b>	WB 1:500-1:2000, IHC-P 1:100-1:300, ELISA 1:10000, IF-P/IF-F/ICC/IF 1:50-200
<b>Molecular Weight</b>	135kDa

## Background


**Product Name: ADAMTS-18 Rabbit Polyclonal Antibody**  
**Catalog #: APRab06599**




This gene encodes a member of the ADAMTS (a disintegrin and metalloproteinase with thrombospondin motifs) protein family. ADAMTS family members share several distinct protein modules, including a propeptide region, a metalloproteinase domain, a disintegrin-like domain, and a thrombospondin type 1 (TS) motif. Individual members of this family differ in the number of C-terminal TS motifs, and some have unique C-terminal domains. The encoded preproprotein is proteolytically processed to generate the mature protein, which may regulate hemostatic balance and function as a tumor suppressor. Mutations in this gene may be associated with microcornea, myopic chorioretinal atrophy, and telecanthus (MMCAT) and cone-rod dystrophy in human patients. [provided by RefSeq, May 2016],cofactor:Binds 1 zinc ion per subunit.,domain:The conserved cysteine present in the cysteine-switch motif binds the catalytic zinc ion, thus inhibiting the enzyme. The dissociation of the cysteine from the zinc ion upon the activation-peptide release activates the enzyme.,PTM:The precursor is cleaved by a furin endopeptidase.,similarity:Contains 1 disintegrin domain.,similarity:Contains 1 peptidase M12B domain.,similarity:Contains 1 PLAC domain.,similarity:Contains 5 TSP type-1 domains.,tissue specificity:Expressed in fetal lung, liver, and kidney and in adult brain, prostate, submaxillary gland, and endothelium.,

## Research Area

## Image Data




Western blot analysis of 3T3 using ADAMTS-18 antibody.. Secondary antibody was diluted at 1:20000



Immunohistochemical analysis of paraffin-embedded human-brain, antibody was diluted at 1:200

**Product Name: ADAMTS-18 Rabbit Polyclonal Antibody**  
**Catalog #: APRab06599**

---



Immunohistochemical analysis of paraffin-embedded human-skin, antibody was diluted at 1:200

### **Note**

For research use only.