# Product Name: Ah Receptor Rabbit Polyclonal Antibody Enkilife Catalog #: APRab06689

# **Summary**

Ah Receptor Rabbit Polyclonal Antibody **Production Name** 

Description Rabbit Polyclonal Antibody

Host Rabbit

**Application** WB,IHC-P,IF-P,IF-F,ICC/IF,ELISA

Reactivity Human, Mouse, Rat

## **Performance**

Conjugation Unconjugated Modification Unmodified

Isotype lgG

Clonality Polyclonal Form Liquid

Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw Storage

cycles.

**Buffer** Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% New type preservative N.

**Purification** Affinity purification

#### **Immunogen**

**Gene Name** AHR

AHR; BHLHE76; Aryl hydrocarbon receptor; Ah receptor; AhR; Class E basic helix-loop-

**Alternative Names** helix protein 76; bHLHe76; AHRR; BHLHE77; KIAA1234; Aryl hydrocarbon receptor

repressor; AhR repressor; AhRR; Class E basic helix-loop-helix protein 77; bHL

Gene ID 196/57491

P35869/A9YTQ3.The antiserum was produced against synthesized peptide derived SwissProt ID

from human AhR. AA range:2-51

# **Application**

**Dilution Ratio** 

WB 1:500-1:2000, IHC-P 1:100-1:300, IF-P/IF-F/ICC/IF 1:200-1:1000, ELISA 1:20000.Not

yet tested in other applications.

Web: https://www.enkilife.com E-mail: order@enkilife.com techsupport@enkilife.com Tel: 0086-27-87002838



**Molecular Weight** 


75 or 96kDa

### Background

The protein encoded by this gene is a ligand-activated helix-loop-helix transcription factor involved in the regulation of biological responses to planar aromatic hydrocarbons. This receptor has been shown to regulate xenobiotic-metabolizing enzymes such as cytochrome P450. Before ligand binding, the encoded protein is sequestered in the cytoplasm; upon ligand binding, this protein moves to the nucleus and stimulates transcription of target genes. [provided by RefSeg, Sep 2015], function: Ligand-activated transcriptional activator. Binds to the XRE promoter region of genes it activates. Activates the expression of multiple phase I and II xenobiotic chemical metabolizing enzyme genes (such as the CYP1A1 gene). Mediates biochemical and toxic effects of halogenated aromatic hydrocarbons. Involved in cell-cycle regulation. Likely to play an important role in the development and maturation of many tissues., induction: Induced or repressed by TGF-beta and dioxin in a cell-type specific fashion. Repressed by cAMP, retinoic acid, and TPA., similarity: Contains 1 basic helix-loophelix (bHLH) domain., similarity: Contains 1 PAC (PAS-associated C-terminal) domain., similarity: Contains 2 PAS (PER-ARNT-SIM) domains., subcellular location: Initially cytoplasmic; upon binding with ligand and interaction with a HSP90, it translocates to the nucleus., subunit: Binds MYBBP1A (By similarity). Efficient DNA binding requires dimerization with another bHLH protein. In the nucleus, heterodimer of AHR and ARNT. Interacts with coactivators including SRC-1, RIP140 and NOCA7, and with the corepressor SMRT. Interacts with NEDD8 and IVNS1ABP., tissue specificity: Expressed in all tissues tested including blood, brain, heart, kidney, liver, lung, pancreas and skeletal muscle.,

#### Research Area


#### **Image Data**




Immunofluorescence analysis of HeLa cells, using AhR Antibody. The picture on the right is blocked with the synthesized peptide.

Web: https://www.enkilife.com E-mail: order@enkilife.com techsupport@enkilife.com Tel: 0086-27-87002838






Immunohistochemistry analysis of paraffin-embedded human breast carcinoma tissue, using AhR Antibody. The picture on the right is blocked with the synthesized peptide.




Western blot analysis of lysates from 293 cells, using AhR Antibody. The lane on the right is blocked with the synthesized peptide.



Western blot analysis of the lysates from HT-29 cells using AhR antibody.







Western Blot analysis of various cells using Ah Receptor Polyclonal Antibody

#### Note

For research use only.