

**Product Name: AID Rabbit Polyclonal Antibody**  
**Catalog #: APRab06697**



## Summary

<b>Production Name</b>	AID Rabbit Polyclonal Antibody
<b>Description</b>	Rabbit Polyclonal Antibody
<b>Host</b>	Rabbit
<b>Application</b>	WB,IHC-P,IF-P,IF-F,ICC/IF,ELISA
<b>Reactivity</b>	Human,Mouse,Rat

## Performance

<b>Conjugation</b>	Unconjugated
<b>Modification</b>	Unmodified
<b>Isotype</b>	IgG
<b>Clonality</b>	Polyclonal
<b>Form</b>	Liquid
<b>Storage</b>	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.
<b>Buffer</b>	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% New type preservative N.
<b>Purification</b>	Affinity purification

## Immunogen

<b>Gene Name</b>	AICDA
<b>Alternative Names</b>	AICDA; AID; Activation-induced cytidine deaminase; Cytidine aminohydrolase
<b>Gene ID</b>	57379.0
<b>SwissProt ID</b>	Q9GZX7.The antiserum was produced against synthesized peptide derived from the Internal region of human AICDA. AA range:81-130

## Application

<b>Dilution Ratio</b>	WB 1:500-1:2000, IHC-P 1:100-1:300, ELISA 1:10000, IF-P/IF-F/ICC/IF 1:50-200
<b>Molecular Weight</b>	24kDa

## Background

**Product Name: AID Rabbit Polyclonal Antibody**  
**Catalog #: APRab06697**




This gene encodes a RNA-editing deaminase that is a member of the cytidine deaminase family. The protein is involved in somatic hypermutation, gene conversion, and class-switch recombination of immunoglobulin genes. Defects in this gene are the cause of autosomal recessive hyper-IgM immunodeficiency syndrome type 2 (HIGM2). [provided by RefSeq, Feb 2009],catalytic activity:Cytidine + H(2)O = uridine + NH(3),cofactor:Zinc,disease:Defects in AICDA are the cause of autosomal recessive hyper-IgM immunodeficiency syndrome type 2 (HIGM2) [MIM:605258]. HIGM2 is characterized by normal or elevated serum IgM levels with absence of IgG, IgA, and IgE, resulting in a profound susceptibility to bacterial infections. HIGM2 causes the absence of Ig class switch recombination (CSR), the lack of Ig somatic hypermutations, and lymph node hyperplasia caused by the presence of giant germinal centers,function:RNA-editing deaminase involved in somatic hypermutation, gene conversion, and class-switch recombination. Required for several crucial steps of B-cell terminal differentiation necessary for efficient antibody responses,online information:AICDA mutation db,similarity:Belongs to the cytidine and deoxycytidylate deaminase family.,tissue specificity:Strongly expressed in lymph nodes and tonsils,.


## Research Area

Intestinal immune network for IgA production;Primary immunodeficiency;

## Image Data



Western Blot analysis of HeLa, HepG2 cells using AID Polyclonal Antibody.. Secondary antibody was diluted at 1:20000



Immunohistochemical analysis of paraffin-embedded human-colon-cancer, antibody was diluted at 1:100

**Product Name: AID Rabbit Polyclonal Antibody**  
**Catalog #: APRab06697**

---



**Note**

For research use only.