

Summary

Production Name	AMPD2 Rabbit Polyclonal Antibody
Description	Rabbit Polyclonal Antibody
Host	Rabbit
Application	WB,ELISA
Reactivity	Human, Mouse, Rat

Performance

Conjugation	Unconjugated
Modification	Unmodified
lsotype	IgG
Clonality	Polyclonal
Form	Liquid
Storage	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw
	cycles.
Buffer	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% New type preservative N.
Purification	Affinity purification

Immunogen

Gene Name	AMPD2
Alternative Names	AMPD2; AMP deaminase 2; AMP deaminase isoform L
Gene ID	271.0
SwissProt ID	Q01433. The antiserum was produced against synthesized peptide derived from human
	AMPD2. AA range:131-180

Application

Dilution Ratio	WB 1:500-1:2000, ELISA 1:5000.Not yet tested in other applications.
Molecular Weight	100kDa

Background

Product Name: AMPD2 Rabbit Polyclonal Antibody Catalog #: APRab06835



The protein encoded by this gene is important in purine metabolism by converting AMP to IMP. The encoded protein, which acts as a homotetramer, is one of three AMP deaminases found in mammals. Several transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, Apr 2012],catalytic activity:AMP + H(2)O = IMP + NH(3).,function:AMP deaminase plays a critical role in energy metabolism.,pathway:Purine metabolism; IMP biosynthesis via salvage pathway; IMP from AMP: step 1/1.,similarity:Belongs to the adenosine and AMP deaminases family.,subunit:Homotetramer.,tissue specificity:Three isoforms are present in mammals: AMP deaminase 1 is the predominant form in skeletal muscle; AMP deaminase 2 predominates in smooth muscle, non-muscle tissue, embryonic muscle and undifferentiated myoblasts; AMP deaminase 3 is found in erythrocytes.,

Research Area

Purine metabolism;

Image Data



Western blot analysis of the lysates from HeLa cells using AMPD2 antibody.



Western Blot analysis of various cells using AMPD2 Polyclonal Antibody

Note

For research use only.