

Summary

Production Name	BOCT Rabbit Polyclonal Antibody
Description	Rabbit Polyclonal Antibody
Host	Rabbit
Application	WB,IHC-P,IF-P,IF-F,ICC/IF,ELISA
Reactivity	Human,Mouse,Rat

Performance

Conjugation	Unconjugated
Modification	Unmodified
lsotype	IgG
Clonality	Polyclonal
Form	Liquid
Storage	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw
	cycles.
Buffer	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% New type preservative N.
Purification	Affinity purification

Immunogen

SLC22A17
SLC22A17; BOCT; BOIT; Solute carrier family 22 member 17; 24p3 receptor; 24p3R;
Brain-type organic cation transporter; Lipocalin-2 receptor; Neutrophil gelatinase-
associated lipocalin receptor; NgalR
51310.0
Q8WUG5.Synthesized peptide derived from the Internal region of human BOCT.

Application

Dilution Ratio	WB 1:500-1:2000, IHC-P 1:100-300, ELISA 1:5000, IF-P/IF-F/ICC/IF 1:50-200
Molecular Weight	57kDa



Background

similarity:Belongs to the major facilitator superfamily. Organic cation transporter family.,tissue specificity:Expressed in brain.,similarity:Belongs to the major facilitator superfamily. Organic cation transporter family.,tissue specificity:Expressed in brain.,

Research Area





Western Blot analysis of extracts from K562, Jurkat cells, using BOCT Polyclonal Antibody.. Secondary antibody was diluted at



Immunohistochemical analysis of paraffin-embedded mouse-brain, antibody was diluted at 1:100





Immunohistochemical analysis of paraffin-embedded mouse-brain, antibody was diluted at 1:100

Note For research use only.