

Product Name: CD298 Rabbit Polyclonal Antibody
Catalog #: APRab08330



Summary

Production Name	CD298 Rabbit Polyclonal Antibody
Description	Rabbit Polyclonal Antibody
Host	Rabbit
Application	WB,ELISA
Reactivity	Human,Rat,Mouse

Performance

Conjugation	Unconjugated
Modification	Unmodified
Isotype	IgG
Clonality	Polyclonal
Form	Liquid
Storage	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.
Buffer	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% New type preservative N.
Purification	Affinity purification

Immunogen

Gene Name	ATP1B3
Alternative Names	ATP1B3; Sodium/potassium-transporting ATPase subunit beta-3; Sodium/potassium-dependent ATPase subunit beta-3; ATPB-3; CD antigen CD298
Gene ID	483.0
SwissProt ID	P54709. Synthesized peptide derived from Human N-terminal CD298 . at AA range: 60-140

Application

Dilution Ratio	WB 1:500-1:2000, ELISA 1:10000.Not yet tested in other applications.
Molecular Weight	31kDa

Background

Product Name: CD298 Rabbit Polyclonal Antibody
Catalog #: AP Rab08330



The protein encoded by this gene belongs to the family of Na⁺/K⁺ and H⁺/K⁺ ATPases beta chain proteins, and to the subfamily of Na⁺/K⁺ -ATPases. Na⁺/K⁺ -ATPase is an integral membrane protein responsible for establishing and maintaining the electrochemical gradients of Na and K ions across the plasma membrane. These gradients are essential for osmoregulation, for sodium-coupled transport of a variety of organic and inorganic molecules, and for electrical excitability of nerve and muscle. This enzyme is composed of two subunits, a large catalytic subunit (alpha) and a smaller glycoprotein subunit (beta). The beta subunit regulates, through assembly of alpha/beta heterodimers, the number of sodium pumps transported to the plasma membrane. The glycoprotein subunit of Na⁺/K⁺ -ATPase is encoded by multiple genes. This gene encodes a beta 3 subunit. This gene encodes a beta 3 subunfunction: This is the non-catalytic component of the active enzyme, which catalyzes the hydrolysis of ATP coupled with the exchange of Na(+) and K(+) ions across the plasma membrane. The exact function of the beta-3 subunit is not known., similarity: Belongs to the X(+)/potassium ATPases subunit beta family., subcellular location: Identified by mass spectrometry in melanosome fractions from stage I to stage IV., subunit: Composed of three subunits: alpha (catalytic), beta and gamma.,

Research Area

Cardiac muscle contraction; Aldosterone-regulated sodium reabsorption;

Image Data



Western Blot analysis of various cells using CD298 Polyclonal Antibody diluted at 1: 1000

Note

For research use only.