

**Product Name: CSN3 Rabbit Polyclonal Antibody**  
**Catalog #: APRab09458**



## Summary

<b>Production Name</b>	CSN3 Rabbit Polyclonal Antibody
<b>Description</b>	Rabbit Polyclonal Antibody
<b>Host</b>	Rabbit
<b>Application</b>	WB,IHC-P,IF-P,IF-F,ICC/IF,ELISA
<b>Reactivity</b>	Human,Mouse,Rat

## Performance

<b>Conjugation</b>	Unconjugated
<b>Modification</b>	Unmodified
<b>Isotype</b>	IgG
<b>Clonality</b>	Polyclonal
<b>Form</b>	Liquid
<b>Storage</b>	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.
<b>Buffer</b>	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% New type preservative N.
<b>Purification</b>	Affinity purification

## Immunogen

<b>Gene Name</b>	COPS3
<b>Alternative Names</b>	COPS3; CSN3; COP9 signalosome complex subunit 3; SGN3; Signalosome subunit 3; JAB1-containing signalosome subunit 3
<b>Gene ID</b>	8533.0
<b>SwissProt ID</b>	Q9UNS2.The antiserum was produced against synthesized peptide derived from human JAB1. AA range:374-423

## Application


<b>Dilution Ratio</b>	WB 1:500-1:2000, IHC-P 1:100-1:300, IF-P/IF-F/ICC/IF 1:200-1:1000, ELISA 1:10000.Not yet tested in other applications.
<b>Molecular Weight</b>	47kDa

## Background

The protein encoded by this gene possesses kinase activity that phosphorylates regulators involved in signal transduction. It phosphorylates I kappa-Balpha, p105, and c-Jun. It acts as a docking site for complex-mediated phosphorylation. The gene is located within the Smith-Magenis syndrome region on chromosome 17. Several transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, Oct 2015],function:Component of the COP9 signalosome complex (CSN), a complex involved in various cellular and developmental processes. The CSN complex is an essential regulator of the ubiquitin (Ubl) conjugation pathway by mediating the deneddylation of the cullin subunits of SCF-type E3 ligase complexes, leading to decrease the Ubl ligase activity of SCF-type complexes such as SCF, CSA or DDB2. The complex is also involved in phosphorylation of p53/TP53, c-jun/JUN, I kappa Balpha/NFKBIA, ITPK1 and IRF8/ICSBP, possibly via its association with CK2 and PKD kinases. CSN-dependent phosphorylation of TP53 and JUN promotes and protects degradation by the Ubl system, respectively.,miscellaneous:Amplified and overexpressed in some osteosarcomas (OS), suggesting that it may participate in TP53 degradation in OS.,PTM:Phosphorylated upon DNA damage, probably by ATM or ATR.,similarity:Belongs to the CSN3 family.,similarity:Contains 1 PCI domain.,subunit:Component of the CSN complex, composed of COPS1/GPS1, COPS2, COPS3, COPS4, COPS5, COPS6, COPS7 (COPS7A or COPS7B) and COPS8. In the complex, it probably interacts directly with COPS1, COPS4 and COPS8. Interacts with CK2 and PKD. Interacts with the translation initiation factor EIF3S6 and IKBKG.,tissue specificity:Widely expressed. Expressed at high level in heart and skeletal muscle.,


## Research Area

## Image Data




Immunofluorescence analysis of HeLa cells, using JAB1 Antibody. The picture on the right is blocked with the synthesized peptide.


**Product Name: CSN3 Rabbit Polyclonal Antibody**  
**Catalog #: APRab09458**



Western blot analysis of lysates from LOVO cells, using JAB1 Antibody. The lane on the right is blocked with the synthesized peptide.



Western Blot analysis of various cells using CSN3 Polyclonal Antibody



Western Blot analysis of 3T3 cells using CSN3 Polyclonal Antibody

## Note

For research use only.