

Summary

Production Name	DHRS4 Rabbit Polyclonal Antibody
Description	Rabbit Polyclonal Antibody
Host	Rabbit
Application	WB,IHC-P,IF-P,IF-F,ICC/IF,ELISA
Reactivity	Human,Rat,Mouse

Performance

Conjugation	Unconjugated
Modification	Unmodified
Isotype	IgG
Clonality	Polyclonal
Form	Liquid
Storage	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.
Buffer	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% New type preservative N.
Purification	Affinity purification

Immunogen

Gene Name	DHRS4 DHRS4; Dehydrogenase/reductase SDR family member 4; NADPH-dependent carbonyl reductase/NADP-retinol dehydrogenase; CR; PHCR; NADPH-dependent retinol dehydrogenase/reductase; NRDR; humNRDR; Peroxisomal short-chain alcohol dehydrogenase; PSCD
Alternative Names	
Gene ID	10901.0
SwissProt ID	Q9BTZ2.The antiserum was produced against synthesized peptide derived from human DHRS4. AA range:191-240

Application

Dilution Ratio	WB 1:500-1:2000, IHC-P 1:100-1:300, ELISA 1:40000, IF-P/IF-F/ICC/IF 1:50-200
Molecular Weight	32-34kDa


Background

catalytic activity: $R-CHOH-R' + NADP(+) = R-CO-R' + NADPH$, function: Reduces all-trans-retinal and 9-cis retinal. Can also catalyze the oxidation of all-trans-retinol with NADP as co-factor, but with much lower efficiency. Reduces alkyl phenyl ketones and alpha-dicarbonyl compounds with aromatic rings, such as pyrimidine-4-aldehyde, 3-benzoylpyridine, 4-benzoylpyridine, menadione and 4-hexanoylpyridine. Has no activity towards aliphatic aldehydes and ketones, miscellaneous: Inhibited by kaempferol, quercetin, genistein and myristic acid, similarity: Belongs to the short-chain dehydrogenases/reductases (SDR) family, subcellular location: Isoform 1 is peroxisomal, while isoform 4 is not, subunit: Homotetramer, tissue specificity: Isoform 1 is predominantly expressed in normal cervix (at protein level). Isoform 4 is expressed in some neoplastic cervical tissues, but not in normal cervix (at protein level). Isoforms 5 and 6 are expressed in a few neoplastic cervical tissues, catalytic activity: $R-CHOH-R' + NADP(+) = R-CO-R' + NADPH$, function: Reduces all-trans-retinal and 9-cis retinal. Can also catalyze the oxidation of all-trans-retinol with NADP as co-factor, but with much lower efficiency. Reduces alkyl phenyl ketones and alpha-dicarbonyl compounds with aromatic rings, such as pyrimidine-4-aldehyde, 3-benzoylpyridine, 4-benzoylpyridine, menadione and 4-hexanoylpyridine. Has no activity towards aliphatic aldehydes and ketones, miscellaneous: Inhibited by kaempferol, quercetin, genistein and myristic acid, similarity: Belongs to the short-chain dehydrogenases/reductases (SDR) family, subcellular location: Isoform 1 is peroxisomal, while isoform 4 is not, subunit: Homotetramer, tissue specificity: Isoform 1 is predominantly expressed in normal cervix (at protein level). Isoform 4 is expressed in some neoplastic cervical tissues, but not in normal cervix (at protein level). Isoforms 5 and 6 are expressed in a few neoplastic cervical tissues,

Research Area


Retinol metabolism;

Image Data




Immunohistochemistry analysis of paraffin-embedded human breast carcinoma tissue, using DHRS4 Antibody. The picture on the right is blocked with the synthesized peptide.

Product Name: DHRS4 Rabbit Polyclonal Antibody
Catalog #: APRab09971



Western blot analysis of lysates from HepG2 cells, using DHRS4 Antibody. The lane on the right is blocked with the synthesized peptide.



Western Blot analysis of various cells using DHRS4 Polyclonal Antibody diluted at 1: 2000

Note

For research use only.