

**Product Name: FGF-1 Rabbit Polyclonal Antibody**  
**Catalog #: APRab10918**



## Summary

<b>Production Name</b>	FGF-1 Rabbit Polyclonal Antibody
<b>Description</b>	Rabbit Polyclonal Antibody
<b>Host</b>	Rabbit
<b>Application</b>	IHC-P,IF-P,IF-F,ICC/IF,ELISA
<b>Reactivity</b>	Human,Mouse,Rat

## Performance

<b>Conjugation</b>	Unconjugated
<b>Modification</b>	Unmodified
<b>Isotype</b>	IgG
<b>Clonality</b>	Polyclonal
<b>Form</b>	Liquid
<b>Storage</b>	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.
<b>Buffer</b>	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% New type preservative N.
<b>Purification</b>	Affinity purification

## Immunogen

<b>Gene Name</b>	FGF1
<b>Alternative Names</b>	FGF1; FGFA; Fibroblast growth factor 1; FGF-1; Acidic fibroblast growth factor; aFGF; Endothelial cell growth factor; ECGF; Heparin-binding growth factor 1; HBGF-1
<b>Gene ID</b>	2246.0
<b>SwissProt ID</b>	P05230.The antiserum was produced against synthesized peptide derived from human FGF-1. AA range:7-56

## Application

<b>Dilution Ratio</b>	IHC-P 1:100-1:300, ELISA 1:10000, IF-P/IF-F/ICC/IF 1:50-200
<b>Molecular Weight</b>	


## Background

The protein encoded by this gene is a member of the fibroblast growth factor (FGF) family. FGF family members possess broad mitogenic and cell survival activities, and are involved in a variety of biological processes, including embryonic development, cell growth, morphogenesis, tissue repair, tumor growth and invasion. This protein functions as a modifier of endothelial cell migration and proliferation, as well as an angiogenic factor. It acts as a mitogen for a variety of mesoderm- and neuroectoderm-derived cells in vitro, thus is thought to be involved in organogenesis. Multiple alternatively spliced variants encoding different isoforms have been described. [provided by RefSeq, Jan 2009],function:The heparin-binding growth factors are angiogenic agents in vivo and are potent mitogens for a variety of cell types in vitro. There are differences in the tissue distribution and concentration of these 2 growth factors.,miscellaneous:This protein binds heparin, although less strongly than does bFGF.,similarity:Belongs to the heparin-binding growth factors family.,subunit:Monomer. Binds FGFR2. Forms a ternary complex containing 2 molecules each of FGFR2 and FGF1 for 1 heparin molecule. Found in a complex with FGFBP1, FGF1 and FGF2. Interacts with FGFBP1.,


## Research Area

MAPK\_ERK\_Growth;MAPK\_G\_Protein;Regulates Actin and Cytoskeleton;Pathways in cancer;Melanoma;

## Image Data



Immunohistochemistry analysis of FGF-1 antibody in paraffin-embedded human brain tissue.



Immunohistochemical analysis of paraffin-embedded Human brain. Antibody was diluted at 1:100 (4°,overnight) . High-

**Product Name: FGF-1 Rabbit Polyclonal Antibody**  
**Catalog #: APRab10918**

---



pressure and temperature Tris-EDTA,pH8.0 was used for antigen retrieval. Negative control (right) obtained from antibody was pre-absorbed by immunogen peptide.

**Note**

For research use only.