Product Name: Glucosidase IIB Rabbit Polyclonal

Antibody

Catalog #: APRab11487



Summary

Production Name Glucosidase IIβ Rabbit Polyclonal Antibody

Description Rabbit Polyclonal Antibody

Host Rabbit

Application WB,IF-P,IF-F,ICC/IF,ELISA

Reactivity Human, Mouse

Performance

ConjugationUnconjugatedModificationUnmodified

Isotype IgG

Clonality Polyclonal Form Liquid

Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw $\bf Storage$

cycles.

Buffer Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% New type preservative N.

Purification Affinity purification

Immunogen

Gene Name PRKCSH

PRKCSH; G19P1; Glucosidase 2 subunit beta; 80K-H protein; Glucosidase II subunit Alternative Names

beta; Protein kinase C substrate 60.1 kDa protein heavy chain; PKCSH

Gene ID 5589.0

P14314.The antiserum was produced against synthesized peptide derived from human SwissProt ID

GLU2B. AA range:81-130

Application

Dilution Ratio

WB 1:500-1:2000, IF-P/IF-F/ICC/IF 1:200-1:1000, ELISA 1:10000.Not yet tested in other

application

applications.

Product Name: Glucosidase IIB Rabbit Polyclonal

Antibody

Catalog #: APRab11487



Molecular Weight


59kDa

Background

This gene encodes the beta-subunit of glucosidase II, an N-linked glycan-processing enzyme in the endoplasmic reticulum. The encoded protein is an acidic phosphoprotein known to be a substrate for protein kinase C. Mutations in this gene have been associated with the autosomal dominant polycystic liver disease. Alternative splicing results in multiple transcript variants. [provided by RefSeq, Jan 2014], disease:Defects in PRKCSH are a cause of polycystic liver disease (PCLD) [MIM:174050]. PCLD is an autosomal dominant disorder and is characterized by the presence of multiple liver cysts of biliary epithelial origin. PCLD is a distinct clinical and genetic entity that can occur independently from autosomal dominant polycystic kidney disease (ADPKD) [MIM:173900], which in a considerable but uncertain proportion of cases is associated with hepatic cysts.,function:Regulatory subunit of glucosidase II.,pathway:Glycan metabolism; N-glycan metabolism,similarity:Contains 1 PRKCSH domain.,similarity:Contains 2 EF-hand domains.,subunit:Heterodimer of a catalytic alpha subunit (GANAB) and a beta subunit (PRKCSH). Binds glycosylated PTPRC.,

Research Area

Image Data



Immunofluorescence analysis of HeLa cells, using GLU2B Antibody. The picture on the right is blocked with the synthesized peptide.


Web: https://www.enkilife.com E-mail: order@enkilife.com techsupport@enkilife.com Tel: 0086-27-87002838

Product Name: Glucosidase IIβ Rabbit Polyclonal


Antibody

Catalog #: APRab11487






Western blot analysis of lysates from HepG2, 293, and HUVEC cells, using GLU2B Antibody. The lane on the right is blocked with the synthesized peptide.



Western blot analysis of the lysates from 293 cells using GLU2B antibody.




Western Blot analysis of various cells using Glucosidase IIB Polyclonal Antibody

Product Name: Glucosidase IIβ Rabbit Polyclonal

Antibody

Catalog #: APRab11487





Western Blot analysis of A549 cells using Glucosidase II β Polyclonal Antibody

Note

For research use only.

Web: https://www.enkilife.com E-mail: order@enkilife.com techsupport@enkilife.com Tel: 0086-27-87002838