

Summary

Production Name	GRB10 Rabbit Polyclonal Antibody
Description	Rabbit Polyclonal Antibody
Host	Rabbit
Application	WB,IHC-P,IF-P,IF-F,ICC/IF,ELISA
Reactivity	Human, Rat, Mouse

Performance

Conjugation	Unconjugated
Modification	Unmodified
lsotype	IgG
Clonality	Polyclonal
Form	Liquid
Storage	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw
	cycles.
Buffer	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% New type preservative N.
Purification	Affinity purification

Immunogen

Gene Name	GRB10
Alternative Names	GRB10; GRBIR; KIAA0207; Growth factor receptor-bound protein 10; GRB10 adapter
	protein; Insulin receptor-binding protein Grb-IR
Gene ID	2887.0
SwissProt ID	Q13322. The antiserum was produced against synthesized peptide derived from human
	GRB10. AA range:33-82

Application

Dilution Patio	WB 1:500-1:2000, IHC-P 1:100-1:300, IF-P/IF-F/ICC/IF 1:200-1:1000, ELISA 1:5000.Not
	yet tested in other applications.
Molecular Weight	67kDa



Background

The product of this gene belongs to a small family of adapter proteins that are known to interact with a number of receptor tyrosine kinases and signaling molecules. This gene encodes a growth factor receptor-binding protein that interacts with insulin receptors and insulin-like growth-factor receptors. Overexpression of some isoforms of the encoded protein inhibits tyrosine kinase activity and results in growth suppression. This gene is imprinted in a highly isoform- and tissue-specific manner, with expression observed from the paternal allele in the brain, and from the maternal allele in the placental trophoblasts. Alternatively spliced transcript variants encoding different isoforms have been identified. [provided by RefSeq, Oct 2010], alternative products: Additional isoforms seem to exist, function: Plays a functional role in insulin and IGF-I signaling. May serve to positively link the insulin and IGF-I receptors to an uncharacterized mitogenic signaling pathway. Interacts with the cytoplasmic domain of the autophosphorylated insulin receptor which is then inhibited. The interaction is mediated by the SH2 domain. Also binds activated platelet-derived growth factor receptor and epidermal growth factor receptor, similarity:Contains 1 SH2 domain., subunit:Interacts with GIGYF1/PERQ1 and GIGYF2/TNRC15, tissue specificity:Highly expressed in skeletal muscle.,

Research Area

Stem cell pathway; Insulin Receptor

Image Data



Immunofluorescence analysis of HepG2 cells, using GRB10 Antibody. The picture on the right is blocked with the synthesized peptide.





Immunohistochemistry analysis of paraffin-embedded human brain tissue, using GRB10 Antibody. The picture on the right is blocked with the synthesized peptide.



-- 26

Western blot analysis of lysates from NIH/3T3 cells, treated with Insulin 0.01U/ml 15 ', using GRB10 Antibody. The lane on the right is blocked with the synthesized peptide.



Western Blot analysis of various cells using GRB10 Polyclonal Antibody diluted at 1: 2000

Product Name: GRB10 Rabbit Polyclonal Antibody Catalog #: APRab11748





Western Blot analysis of 3T3 cells using GRB10 Polyclonal Antibody diluted at 1: 2000

Note

For research use only.