

Summary

Production Name	IL-6R α Rabbit Polyclonal Antibody
Description	Rabbit Polyclonal Antibody
Host	Rabbit
Application	WB,IHC-P,IF-P,IF-F,ICC/IF,ELISA
Reactivity	Human,Rat,Mouse

Performance

Conjugation	Unconjugated
Modification	Unmodified
lsotype	IgG
Clonality	Polyclonal
Form	Liquid
Storage	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw
	cycles.
Buffer	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% New type preservative N.
Purification	Affinity purification

Immunogen

Gene Name	IL6R
Alternative Names	IL6R; Interleukin-6 receptor subunit alpha; IL-6 receptor subunit alpha; IL-6R subunit
	alpha; IL-6R-alpha; IL-6RA; IL-6R 1; Membrane glycoprotein 80; gp80; CD126
Gene ID	3570.0
SwissProt ID	P08887.The antiserum was produced against synthesized peptide derived from the
	Internal region of human IL6R. AA range:221-270

Application

Dilution Ratio	IHC-P 100-300.WB 1:500-1:2000, ELISA 1:10000, IF-P/IF-F/ICC/IF 1:50-200
Molecular Weight	53kDa




Background

This gene encodes a subunit of the interleukin 6 (IL6) receptor complex. Interleukin 6 is a potent pleiotropic cytokine that regulates cell growth and differentiation and plays an important role in the immune response. The IL6 receptor is a protein complex consisting of this protein and interleukin 6 signal transducer (IL6ST/GP130/IL6-beta), a receptor subunit also shared by many other cytokines. Dysregulated production of IL6 and this receptor are implicated in the pathogenesis of many diseases, such as multiple myeloma, autoimmune diseases and prostate cancer. Alternatively spliced transcript variants encoding distinct isoforms have been reported. A pseudogene of this gene is found on chromosome 9.[provided by RefSeq, May 2011], domain: The two fibronectin type-III-like domains, contained in the N-terminal part, form together a cytokine-binding domain.,domain:The WSXWS motif appears to be necessary for proper protein folding and thereby efficient intracellular transport and cell-surface receptor binding, function:Low concentration of a soluble form of IL6 receptor acts as an agonist of IL6 activity., function: Part of the receptor for interleukin 6. Binds to IL6 with low affinity, but does not transduce a signal. Signal activation necessitate an association with IL6ST. Activation may lead to the regulation of the immune response, acute-phase reactions and hematopoiesis., PTM:A short soluble form may also be released from the membrane by proteolysis, similarity: Belongs to the type I cytokine receptor family. Type 3 subfamily, similarity: Contains 1 fibronectin type-III domain., similarity: Contains 1 Ig-like C2-type (immunoglobulin-like) domain., subunit: Hexamer of two molecules each of IL6, IL6R and IL6ST., tissue specificity: Isoform 2 is expressed in peripheral blood mononuclear cells and weakly found in urine and serum.,

Research Area


Cytokine-cytokine receptor interaction;Jak_STAT;Hematopoietic cell lineage;

Image Data




Western Blot analysis of AD293, 4T1 cells using IL-6Ra Polyclonal Antibody.. Secondary antibody was diluted at 1:20000






Immunohistochemical analysis of paraffin-embedded Human spleen. 1, Antibody was diluted at 1:200 (4°,overnight) . 2, High-pressure and temperature EDTA, pH8.0 was used for antigen retrieval. 3,Secondary antibody was diluted at



1:200 (room temperature, 30min) .

Immunohistochemical analysis of paraffin-embedded Human spleen. 1, Antibody was diluted at 1:200 (4°,overnight) . 2, High-pressure and temperature EDTA, pH8.0 was used for antigen retrieval. 3,Secondary antibody was diluted at



Immunohistochemical analysis of paraffin-embedded Human spleen. 1, Antibody was diluted at 1:200 (4°,overnight) . 2, High-pressure and temperature EDTA, pH8.0 was used for antigen retrieval. 3,Secondary antibody was diluted at 1:200 (room temperature, 30min) . Product Name: IL-6Rα Rabbit Polyclonal Antibody Catalog #: APRab12565



Note

For research use only.