Product Name: NDUFA8 Rabbit Polyclonal Antibody

Catalog #: APRab14500



Summary

Production Name NDUFA8 Rabbit Polyclonal Antibody

Description Rabbit Polyclonal Antibody

Host Rabbit

Application WB,IHC-P,IF-P,IF-F,ICC/IF,ELISA

Reactivity Human, Mouse

Performance

ConjugationUnconjugatedModificationUnmodified

Isotype IgG

ClonalityPolyclonalFormLiquid

Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw

cycles.

Buffer Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% New type preservative N.

Purification Affinity purification

Immunogen

Storage

Gene Name NDUFA8

NDUFA8; NADH dehydrogenase [ubiquinone] 1 alpha subcomplex subunit 8; Complex

Alternative Names I-19kD; CI-19kD; Complex I-PGIV; CI-PGIV; NADH-ubiquinone oxidoreductase 19 kDa

subunit

Gene ID 4702.0

P51970.The antiserum was produced against synthesized peptide derived from human SwissProt ID

NDUFA8. AA range:109-158

Application

Dilution Ratio WB 1:500-1:2000, IHC-P 1:100-1:300, ELISA 1:40000, IF-P/IF-F/ICC/IF 1:50-200

Molecular Weight 22kDa

Product Name: NDUFA8 Rabbit Polyclonal Antibody

Catalog #: APRab14500




Background

The protein encoded by this gene belongs to the complex I 19 kDa subunit family. Mammalian complex I is composed of 45 different subunits. This protein has NADH dehydrogenase activity and oxidoreductase activity. It plays an important role in transfering electrons from NADH to the respiratory chain. The immediate electron acceptor for the enzyme is believed to be ubiquinone. Alternative splicing of this gene results in multiple transcript variants encoding different isoforms. [provided by RefSeq, Dec 2015], function: Accessory subunit of the mitochondrial membrane respiratory chain NADH dehydrogenase (Complex I), that is believed to be not involved in catalysis. Complex I functions in the transfer of electrons from NADH to the respiratory chain. The immediate electron acceptor for the enzyme is believed to be ubiquinone., similarity: Belongs to the complex I NDUFA8 subunit family., similarity: Contains 2 CHCH domains., subunit: Complex I is composed of 45 different subunits..


Research Area

Oxidative phosphorylation; Alzheimer's disease; Parkinson's disease; Huntington's disease;

Image Data



Immunohistochemistry analysis of paraffin-embedded human pancreas tissue, using NDUFA8 Antibody. The picture on the right is blocked with the synthesized peptide.




Western blot analysis of lysates from HUVEC, MCF-7, Jurkat, HepG2, and 293 cells, using NDUFA8 Antibody. The lane on the

Product Name: NDUFA8 Rabbit Polyclonal Antibody Catalog #: APRab14500

G EnkiLife

right is blocked with the synthesized peptide.



Western Blot analysis of various cells using NDUFA8 Polyclonal Antibody diluted at 1: 1000

Note

For research use only.