

Summary

Production Name	NOS3 Rabbit Polyclonal Antibody
Description	Rabbit Polyclonal Antibody
Host	Rabbit
Application	WB,IHC-P,IF-P,IF-F,ICC/IF,ELISA
Reactivity	Human,Mouse,Rat

Performance

Conjugation	Unconjugated
Modification	Unmodified
Isotype	IgG
Clonality	Polyclonal
Form	Liquid
Storage	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.
Buffer	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% New type preservative N.
Purification	Affinity purification

Immunogen

Gene Name	NOS3
Alternative Names	NOS3; Nitric oxide synthase; endothelial; Constitutive NOS; cNOS; EC-NOS; Endothelial NOS; eNOS; NOS type III; NOSIII
Gene ID	4846.0
SwissProt ID	P29474.The antiserum was produced against synthesized peptide derived from human eNOS. AA range:1145-1194

Application

Dilution Ratio	WB 1:500-2000, IF-P/IF-F/ICC/IF 1:50-300, IHC-P 1:50-300
Molecular Weight	130-140kDa


Background

Nitric oxide is a reactive free radical which acts as a biologic mediator in several processes, including neurotransmission and antimicrobial and antitumoral activities. Nitric oxide is synthesized from L-arginine by nitric oxide synthases. Variations in this gene are associated with susceptibility to coronary spasm. Multiple transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, May 2009], catalytic activity: L-arginine + n NADPH + n H⁽⁺⁾ + m O₂ = citrulline + nitric oxide + n NADP⁽⁺⁾., cofactor: Binds 1 FAD., cofactor: Binds 1 FMN., cofactor: Heme group., cofactor: Tetrahydrobiopterin (BH₄). May stabilize the dimeric form of the enzyme., enzyme regulation: Stimulated by calcium/calmodulin. Inhibited by NOSIP and NOSTRIN., function: Produces nitric oxide (NO) which is implicated in vascular smooth muscle relaxation through a cGMP-mediated signal transduction pathway. NO mediates vascular endothelial growth factor (VEGF)-induced angiogenesis in coronary vessels and promotes blood clotting through the activation of platelets., online information: Nitric oxide synthase entry, polymorphism: Variation in NOS3 seem to be associated with susceptibility to coronary spasm., similarity: Belongs to the NOS family., similarity: Contains 1 FAD-binding FR-type domain., similarity: Contains 1 flavodoxin-like domain., subcellular location: Specifically associates with actin cytoskeleton in the G2 phase of the cell cycle; which is favored by interaction with NOSIP and results in a reduced enzymatic activity., subunit: Homodimer. Interacts with NOSIP and NOSTRIN., tissue specificity: Platelets, placenta, liver and kidney.,

Research Area


Regulates Angiogenesis; AMPK; PI3K/Akt; Protein_Acetylation

Image Data




Immunofluorescence analysis of HeLa cells, using eNOS Antibody. The picture on the right is blocked with the synthesized peptide.


Product Name: NOS3 Rabbit Polyclonal Antibody
Catalog #: APRab14804



Western blot analysis of lysates from A549, K562, and COLO205 cells, using eNOS Antibody. The lane on the right is blocked with the synthesized peptide.




Western blot analysis of the lysates from HepG2 cells using eNOS antibody.




Immunofluorescence analysis of rat-lung tissue. 1, NOS3 Polyclonal Antibody (red) was diluted at 1:200 (4°C, overnight).
 2, Cy3 labeled Secondary antibody was diluted at 1:300 (room temperature, 50min). 3, Picture B: DAPI (blue) 10min.
 Picture A: Target. Picture B: DAPI. Picture C: merge of A+B

Product Name: NOS3 Rabbit Polyclonal Antibody
Catalog #: APRab14804




Immunofluorescence analysis of rat-lung tissue. 1, NOS3 Polyclonal Antibody (red) was diluted at 1:200 (4°C, overnight).
2, Cy3 labeled Secondary antibody was diluted at 1:300 (room temperature, 50min). 3, Picture B: DAPI (blue) 10min.

Picture A: Target. Picture B: DAPI. Picture C: merge of A+B



Immunofluorescence analysis of rat-kidney tissue. 1, NOS3 Polyclonal Antibody (red) was diluted at 1:200 (4°C, overnight).
2, Cy3 labeled Secondary antibody was diluted at 1:300 (room temperature, 50min). 3, Picture B: DAPI (blue) 10min.


Picture A: Target. Picture B: DAPI. Picture C: merge of A+B



Immunofluorescence analysis of rat-kidney tissue. 1, NOS3 Polyclonal Antibody (red) was diluted at 1:200 (4°C, overnight).
2, Cy3 labeled Secondary antibody was diluted at 1:300 (room temperature, 50min). 3, Picture B: DAPI (blue) 10min.


Picture A: Target. Picture B: DAPI. Picture C: merge of A+B

Product Name: NOS3 Rabbit Polyclonal Antibody
Catalog #: APRab14804



Immunofluorescence analysis of rat-spleen tissue. 1, NOS3 Polyclonal Antibody (red) was diluted at 1:200 (4°C, overnight) .
2, Cy3 labeled Secondary antibody was diluted at 1:300 (room temperature, 50min) .3, Picture B: DAPI (blue) 10min.

Picture A: Target. Picture B: DAPI. Picture C: merge of A+B



Immunofluorescence analysis of rat-spleen tissue. 1, NOS3 Polyclonal Antibody (red) was diluted at 1:200 (4°C, overnight) .
2, Cy3 labeled Secondary antibody was diluted at 1:300 (room temperature, 50min) .3, Picture B: DAPI (blue) 10min.

Picture A: Target. Picture B: DAPI. Picture C: merge of A+B

Note

For research use only.