

# Summary

Production Name	Olfactory receptor 3A2/3A3 Rabbit Polyclonal Antibody
Description	Rabbit Polyclonal Antibody
Host	Rabbit
Application	WB,IF-P,IF-F,ICC/IF,ELISA
Reactivity	Human, Rat, Mouse

### Performance

Conjugation	Unconjugated
Modification	Unmodified
lsotype	lgG
Clonality	Polyclonal
Form	Liquid
Storage	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw
	cycles.
Buffer	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% New type preservative N.
Purification	Affinity purification

### Immunogen

Gene Name	OR3A2/OR3A3
Alternative Names	OR3A3; OR3A6; OR3A7; OR3A8P; Olfactory receptor 3A3; Olfactory receptor 17-201;
	OR17-201; Olfactory receptor 3A6; Olfactory receptor 3A7; Olfactory receptor 3A8;
	Olfactory receptor OR17-22; OR3A2; OLFRA04; Olfactory receptor 3A2; Olfactory
Gene ID	8392/4995
SwissProt ID	P47888/P47893.The antiserum was produced against synthesized peptide derived from
	human OR3A2/3. AA range:65-114

# Application

<b>Dilution Ratio</b>	WB 1:500-1:2000, IF-P/IF-F/ICC/IF 1:200-1:1000, ELISA 1:5000.Not yet tested in other
-----------------------	--

# Product Name: Olfactory receptor 3A2/3A3 Rabbit Polyclonal Antibody Catalog #: APRab15212



applications.

**Molecular Weight** 

36kDa


# Background

olfactory receptor family 3 subfamily A member 3(OR3A3) Homo sapiens Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptor proteins are members of a large family of G-protein-coupled receptors (GPCR) arising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor gene family is the largest in the genome. The nomenclature assigned to the olfactory receptor genes and proteins for this organism is independent of other organisms. [provided by RefSeq, Jul 2008],caution: It is uncertain whether Met-1 or Met-7 is the initiator, function: Odorant receptor .,similarity: Belongs to the G-protein coupled receptor 1 family.,

# **Research Area**


Olfactory transduction;

# Image Data




Immunofluorescence analysis of MCF7 cells, using OR3A2/3 Antibody. The picture on the right is blocked with the synthesized peptide.





Western blot analysis of lysates from MCF-7 cells, using OR3A2/3 Antibody. The lane on the right is blocked with the





Note

For research use only.