

Summary

Production Name	PA26 Rabbit Polyclonal Antibody
Description	Rabbit Polyclonal Antibody
Host	Rabbit
Application	WB,IHC-P,IF-P,IF-F,ICC/IF,ELISA
Reactivity	Human,Mouse,Rat

Performance

Conjugation	Unconjugated
Modification	Unmodified
lsotype	IgG
Clonality	Polyclonal
Form	Liquid
Storage	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw
	cycles.
Buffer	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% New type preservative N.
Purification	Affinity purification

Immunogen

Gene Name	SESN1
Alternative Names	SESN1; PA26; SEST1; Sestrin-1; p53-regulated protein PA26
Gene ID	27244.0
SwissProt ID	Q9Y6P5.The antiserum was produced against synthesized peptide derived from human
	SESN1. AA range:271-320

Application

Dilution Ratio	WB 1:500-1:2000, IHC-P 1:100-1:300, ELISA 1:20000, IF-P/IF-F/ICC/IF 1:50-200
Molecular Weight	57kDa

Background

Product Name: PA26 Rabbit Polyclonal Antibody Catalog #: APRab15675




This gene encodes a member of the sestrin family. Sestrins are induced by the p53 tumor suppressor protein and play a role in the cellular response to DNA damage and oxidative stress. The encoded protein mediates p53 inhibition of cell growth by activating AMP-activated protein kinase, which results in the inhibition of the mammalian target of rapamycin protein. The encoded protein also plays a critical role in antioxidant defense by regenerating overoxidized peroxiredoxins, and the expression of this gene is a potential marker for exposure to radiation. Alternatively spliced transcript variants encoding multiple isoforms have been observed for this gene. [provided by RefSeq, Dec 2010],disease:Defects in SESN1 may be involved in heterotaxia. Heterotaxia is an aetiologically heterogeneous condition caused by an abnormal left-right axis formation, resulting in reversed left-right polarity of one or more organ systems.,function:Involved in the reduction of peroxiredoxins. May also be regulator of cellular growth.,induction:Isoforms T2 and isoform T3 are induced by genotoxic stress (UV, gamma-irradiation and cytotoxic drugs) in a p53-dependent manner. Isoform T1 is not induced by p53,,PTM:Phosphorylated upon DNA damage, probably by ATM or ATR.,similarity:Belongs to the sestrin family,,tissue specificity:Widely expressed.,


Research Area

p53;

Image Data



Immunohistochemistry analysis of paraffin-embedded human breast carcinoma tissue, using SESN1 Antibody. The picture on



Product Name: PA26 Rabbit Polyclonal Antibody Catalog #: APRab15675



Western blot analysis of lysates from rat muscle cells, using SESN1 Antibody. The lane on the right is blocked with the synthesized peptide.

Note For research use only.