

Product Name: PI 3-Kinase p110δ Rabbit Polyclonal Antibody
Catalog #: APRab16100

Summary

Production Name	PI 3-Kinase p110δ Rabbit Polyclonal Antibody
Description	Rabbit Polyclonal Antibody
Host	Rabbit
Application	WB,IHC-P,IF-P,IF-F,ICC/IF,ELISA
Reactivity	Human,Mouse

Performance

Conjugation	Unconjugated
Modification	Unmodified
Isotype	IgG
Clonality	Polyclonal
Form	Liquid
Storage	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.
Buffer	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% New type preservative N.
Purification	Affinity purification

Immunogen

Gene Name	PIK3CD PIK3CD; Phosphatidylinositol 4,5-bisphosphate 3-kinase catalytic subunit delta isoform; PI3-kinase subunit delta; PI3K-delta; PI3Kdelta; PtdIns-3-kinase subunit delta; Phosphatidylinositol 4,5-bisphosphate 3-kinase 110 kDa catalytic subunit delta; PtdIns-3-kinase subunit p110-delta; p110delta
Alternative Names	
Gene ID	5293.0
SwissProt ID	O00329.The antiserum was produced against synthesized peptide derived from the N-terminal region of human PIK3CD. AA range:41-90

Application

Dilution Ratio	WB 1:500-1:2000, IHC-P 1:100-1:300, ELISA 1:20000, IF-P/IF-F/ICC/IF 1:50-200
-----------------------	--

Product Name: PI 3-Kinase p110δ Rabbit Polyclonal Antibody
Catalog #: AP Rab16100



Molecular Weight 120kDa


Background

Phosphoinositide 3-kinases (PI3Ks) phosphorylate inositol lipids and are involved in the immune response. The protein encoded by this gene is a class I PI3K found primarily in leukocytes. Like other class I PI3Ks (p110-alpha p110-beta, and p110-gamma), the encoded protein binds p85 adapter proteins and GTP-bound RAS. However, unlike the other class I PI3Ks, this protein phosphorylates itself, not p85 protein.[provided by RefSeq, Jul 2010],catalytic activity:ATP + 1-phosphatidyl-1D-myo-inositol 4,5-bisphosphate = ADP + 1-phosphatidyl-1D-myo-inositol 3,4,5-trisphosphate.,pathway:Phospholipid metabolism; phosphatidylinositol phosphate biosynthesis.,PTM:Autophosphorylation on Ser-1039 results in the almost complete inactivation of the lipid kinase activity.,similarity:Belongs to the PI3/PI4-kinase family.,similarity:Contains 1 PI3K/PI4K domain.,subunit:Heterodimer of a p110 (catalytic) and a p85 (regulatory) subunit. Interacts with ERAS.,tissue specificity:Expressed predominantly in leukocytes.,

Research Area


Inositol phosphate metabolism;ErbB_HER;Chemokine;Phosphatidylinositol signaling system;mTOR;Apoptosis_Inhibition;Apoptosis_Mitochondrial;Apoptosis_Overview;VEGF;Focal adhesion;Toll_Like;Jak_STAT;Natural killer cell mediated cytotoxicity;T_Cell_Receptor;B_Cell_Antigen;Fc epsilon RI;Fc gamma R-mediated phagocytosis;Leukocyte transendothelial migration;Neurotrophin;Regulates Actin and Cytoskeleton;Insulin_Receptor;Progesterone-mediated oocyte maturation;Type II diabetes mellitus;Aldosterone-regulated sodium reabsorption;Pathways in cancer;Colorectal cancer;Renal cell carcinoma;Pancreatic cancer;Endometrial cancer;Glioma;Prostate cancer;Melanoma;Chronic myeloid leukemia;Acute myeloid leukemia;Small cell lung cancer;Non-small cell lung cancer;

Image Data



Western Blot analysis of HepG2 cells using PI 3-Kinase p110δ Polyclonal Antibody.. Secondary antibody was diluted at 1:20000

**Product Name: PI 3-Kinase p110δ Rabbit Polyclonal
Antibody**
Catalog #: APRab16100



Immunohistochemical analysis of paraffin-embedded mouse-brain, antibody was diluted at 1:100

Note

For research use only.