

Product Name: S2P Rabbit Polyclonal Antibody
Catalog #: APRab17522



Summary

Production Name	S2P Rabbit Polyclonal Antibody
Description	Rabbit Polyclonal Antibody
Host	Rabbit
Application	WB,IP,IF-P,IF-F,ICC/IF,ELISA
Reactivity	Human,Mouse,Rat

Performance

Conjugation	Unconjugated
Modification	Unmodified
Isotype	IgG
Clonality	Polyclonal
Form	Liquid
Storage	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.
Buffer	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% New type preservative N.
Purification	Affinity purification

Immunogen

Gene Name	MBTPS2 MBTPS2; S2P; Membrane-bound transcription factor site-2 protease; Endopeptidase
Alternative Names	S2P; Sterol regulatory element-binding proteins intramembrane protease; SREBPs intramembrane protease
Gene ID	51360.0
SwissProt ID	O43462. The antiserum was produced against synthesized peptide derived from human MBTPS2. AA range:301-350

Application

Dilution Ratio	WB 1:500-2000, IF-P/IF-F/ICC/IF 1:200-1:1000, ELISA 1:20000. Not yet tested in other applications.
Molecular Weight	

Product Name: S2P Rabbit Polyclonal Antibody
Catalog #: APRab17522




Background

This gene encodes a intramembrane zinc metalloprotease, which is essential in development. This protease functions in the signal protein activation involved in sterol control of transcription and the ER stress response. Mutations in this gene have been associated with ichthyosis follicularis with atrichia and photophobia (IFAP syndrome); IFAP syndrome has been quantitatively linked to a reduction in cholesterol homeostasis and ER stress response.[provided by RefSeq, Aug 2009],catalytic activity: Cleaves several transcription factors that are type-2 transmembrane proteins within membrane-spanning domains. Known substrates include sterol regulatory element-binding protein (SREBP) -1, SREBP-2 and forms of the transcriptional activator ATF6. SREBP-2 is cleaved at the site 477-DRSRILL-[-CVLTFLCLSFNPLTSLLQWGGA-505. The residues Asn-Pro, 11 residues distal to the site of cleavage in the membrane-spanning domain, are important for cleavage by S2P endopeptidase. Replacement of either of these residues does not prevent cleavage, but there is no cleavage if both of these residues are replaced.,cofactor: Binds 1 zinc ion per subunit.,function: Intramembrane proteolysis of sterol-regulatory element-binding proteins (SREBPs) within the first transmembrane segment thereby releasing the N-terminal segment with a portion of the transmembrane segment attached. Site-2 cleavage comes after site-1 cleavage which takes place in the luminal loop.,similarity: Belongs to the peptidase M50A family.,tissue specificity: Expressed in heart, brain, placenta, lung, liver, muscle, kidney and pancreas.,

Research Area

Image Data



Immunofluorescence analysis of A549 cells, using MBTPS2 Antibody. The picture on the right is blocked with the synthesized peptide.

Note

For research use only.