

## Summary

|                        |                                   |
|------------------------|-----------------------------------|
| <b>Production Name</b> | TIF-IA Rabbit Polyclonal Antibody |
| <b>Description</b>     | Rabbit Polyclonal Antibody        |
| <b>Host</b>            | Rabbit                            |
| <b>Application</b>     | WB,IHC-P                          |
| <b>Reactivity</b>      | Human,Mouse,Rat                   |

## Performance

|                     |  |
|---------------------|--|
| <b>Conjugation</b>  | Unconjugated   |
| <b>Modification</b> | Unmodified   |
| <b>Isotype</b>      | IgG  |
| <b>Clonality</b>    | Polyclonal   |
| <b>Form</b>         | Liquid   |
| <b>Storage</b>      | Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles. |
| <b>Buffer</b>       | Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% New type preservative N.       |
| <b>Purification</b> | Affinity purification  |

## Immunogen

|                          |  |
|--------------------------|--|
| <b>Gene Name</b>         | RRN3 TIFIA   |
| <b>Alternative Names</b> | RNA polymerase I-specific transcription initiation factor RRN3 (Transcription initiation factor IA) (TIF-IA) |
| <b>Gene ID</b>           | 54700.0  |
| <b>SwissProt ID</b>      | Q9NYV6. Synthesized peptide derived from human TIF-IA Polyclonal   |

## Application

|                         |                               |
|-------------------------|-------------------------------|
| <b>Dilution Ratio</b>   | WB 1:500-2000, IHC-P 1:50-300 |
| <b>Molecular Weight</b> | 70kDa                         |

## Background

function:Required for efficient transcription initiation by RNA polymerase I.,similarity:Belongs to the RRN3


**Product Name: TIF-IA Rabbit Polyclonal Antibody**  
**Catalog #: APRab18930**



family, function: Required for efficient transcription initiation by RNA polymerase I., similarity: Belongs to the RRN3 family.,

## Research Area

## Image Data



Western blot analysis of 3T3 lysate, antibody was diluted at 1000. Secondary antibody was diluted at 1:20000

## Note

For research use only.