

Summary

| | |
|------------------------|------------------------------------|
| Production Name | TudorSN Rabbit Polyclonal Antibody |
| Description | Rabbit Polyclonal Antibody |
| Host | Rabbit |
| Application | WB,IHC-P,IF-P,IF-F,ICC/IF,ELISA |
| Reactivity | Human,Mouse,Rat |

Performance

| | |
|---------------------|--|
| Conjugation | Unconjugated |
| Modification | Unmodified |
| Isotype | IgG |
| Clonality | Polyclonal |
| Form | Liquid |
| Storage | Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles. |
| Buffer | Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% New type preservative N. |
| Purification | Affinity purification |

Immunogen

| | |
|--------------------------|---|
| Gene Name | SND1 |
| Alternative Names | SND1; TDRD11; Staphylococcal nuclease domain-containing protein 1; 100 kDa coactivator; EBNA2 coactivator p100; Tudor domain-containing protein 11; p100 co-activator |
| Gene ID | 27044.0 |
| SwissProt ID | Q7KZF4.Synthesized peptide derived from the Internal region of human TudorSN. |

Application


| | |
|-------------------------|--|
| Dilution Ratio | WB 1:500-1:2000, IHC-P 1:100-300, ELISA 1:40000, IF-P/IF-F/ICC/IF 1:50-200 |
| Molecular Weight | 101kDa |

Background


This gene encodes a transcriptional co-activator that interacts with the acidic domain of Epstein-Barr virus nuclear antigen 2 (EBNA 2), a transcriptional activator that is required for B-lymphocyte transformation. Other transcription factors that interact with this protein are signal transducers and activators of transcription, STATs. This protein is also thought to be essential for normal cell growth. A similar protein in mammals and other organisms is a component of the RNA-induced silencing complex (RISC). [provided by RefSeq, Jul 2016],function:Functions as a bridging factor between STAT6 and the basal transcription factor. Plays a role in PIM1 regulation of MYB activity. Functions as a transcriptional coactivator for the Epstein-Barr virus nuclear antigen 2 (EBNA2),PTM:Phosphorylated by PIM1 in vitro.,sequence caution:The frameshift leads to wrong initiation.,similarity:Contains 1 Tudor domain.,similarity:Contains 4 TNase-like domains.,subcellular location:In IL-4 stimulated cells colocalizes with STAT6 in the nucleus. Identified by mass spectrometry in melanosome fractions from stage I to stage IV.,subunit:Binds to acidic transactivation domain of EBNA2. Interacts with EAV NSP1. Interacts with GTF2E1 and GTF2E2. Forms a ternary complex with STAT6 and POLR2A. Interacts with STAT5.,tissue specificity:Ubiquitously expressed.,

Research Area

Image Data




Western Blot analysis of extracts from Jurkat cells, using TudorSN Polyclonal Antibody.. Secondary antibody was diluted at 1:20000




Product Name: TudorSN Rabbit Polyclonal Antibody
Catalog #: APRab19429



Immunohistochemical analysis of paraffin-embedded mouse-brain, antibody was diluted at 1:100



Immunohistochemical analysis of paraffin-embedded mouse-brain, antibody was diluted at 1:100



Immunohistochemical analysis of paraffin-embedded mouse-brain, antibody was diluted at 1:100

Note

For research use only.