

Summary

| Production Name | β -1,4-GalNAc-T Rabbit Polyclonal Antibody |
|-----------------|--|
| Description | Rabbit Polyclonal Antibody |
| Host | Rabbit |
| Application | WB,IHC-P,IF-P,IF-F,ICC/IF,ELISA |
| Reactivity | Human,Mouse,Rat |
| | |

Performance

| Conjugation | Unconjugated |
|--------------|--|
| Modification | Unmodified |
| lsotype | lgG |
| Clonality | Polyclonal |
| Form | Liquid |
| Storage | Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles. |
| Buffer | Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% New type preservative N. |
| Purification | Affinity purification |

Immunogen

| Gene Name | CSGALNACT1 | | | | | | |
|-------------------|---|-----------------|-------------------|---------|------|--|--|
| | CSGALNACT1; CHGN; | GALNACT1; | Chondroitin | sulfate | N- | | |
| Alternative Names | acetylgalactosaminyltransfera | ise 1; CsGalNAd | cT-1; Chondroitin | beta-1; | 4-N- | | |
| | acetylgalactosaminyltransferase 1; Beta4GalNAcT-1 | | | | | | |
| Gene ID | 55790.0 | | | | | | |
| SwissProt ID | Q8TDX6. The antiserum was produced against synthesized peptide derived from | | | | | | |
| | human CSGALNACT1. AA ran | ge:201-250 | | | | | |

Application

| Dilution Ratio | WB 1:500-1:2000, IHC-P 1:100-1:300, ELISA 1:40000, IF-P/IF-F/ICC/IF 1:50-200 |
|------------------|--|
| Molecular Weight | 60kDa |



Background

catalytic activity:UDP-N-acetyl-D-galactosamine + beta-D-glucuronyl-(1->3)-D-galactosyl-proteoglycan = UDP + N-acetyl-D-galactosaminyl-(1->4)-beta-D-glucuronyl-(1->3)-beta-D-galactosylproteoglycan.,function:Transfers 1,4-Nacetylgalactosamine (GalNAc) from UDP-GalNAc to the non-reducing end of glucuronic acid (GlcUA). Required for addition of the first GalNAc to the core tetrasaccharide linker and for elongation of chondroitin chains. Important role in chondroitin chain biosynthesis in cartilage., online information: Chondroitin beta-1,4-N-acetylgalactosaminyltransferase 1, online information:GlycoGene database,PTM:N-glycosylated.,similarity:Belongs to the chondroitin Nacetylgalactosaminyltransferase family.,tissue specificity:Ubiquitous, with the highest levels in placenta, thyroid, bladder, prostate and adrenal gland. Detected at low levels in the other tissues examined, catalytic activity:UDP-N-acetyl-Dgalactosamine + beta-D-glucuronyl-(1->3)-D-galactosyl-proteoglycan = UDP + N-acetyl-D-galactosaminyl-(1->4)-beta-Dglucuronyl-(1->3)-beta-D-galactosylproteoglycan.,function:Transfers 1,4-N-acetylgalactosamine (GalNAc) from UDP-GalNAc to the non-reducing end of glucuronic acid (GlcUA). Required for addition of the first GalNAc to the core tetrasaccharide linker and for elongation of chondroitin chains. Important role in chondroitin chain biosynthesis in cartilage.,online information:Chondroitin beta-1,4-N-acetylgalactosaminyltransferase 1,online information:GlycoGene database, PTM:N-glycosylated., similarity: Belongs to the chondroitin N-acetylgalactosaminyltransferase family., tissue specificity:Ubiquitous, with the highest levels in placenta, thyroid, bladder, prostate and adrenal gland. Detected at low levels in the other tissues examined.,

Research Area

Chondroitin sulfate biosynthesis;

Image Data



Immunohistochemistry analysis of paraffin-embedded human breast carcinoma tissue, using CSGALNACT1 Antibody. The picture on the right is blocked with the synthesized peptide.



Note

For research use only.