

Product Name: KD-Validated Pro Caspase 3 Recombinant Rabbit Monoclonal Antibody
Catalog #: KVA00230

For research use only.

Summary

Description	KO&KD-Validated antibody
Host	Rabbit
Application	WB,FCM,IHC-P
Reactivity	Human
Conjugation	Unconjugated
Modification	Unmodified
Isotype	Rabbit IgG
Clonality	Rabbit mAb
Form	Liquid
Storage	Aliquot and store at -20°C (valid for 12 months). Avoid freeze/thaw cycles.
Shipping	Ice bags
Buffer	Supplied in PBS (pH 7.4) containing 50% glycerol, and 0.02% sodium azide.
Purification	Affinity purification

Application

Dilution Ratio	WB 1:1,000-1:5,000; FC 1:200-1:2,000; IHC-P 1:100-1:200
Molecular Weight	Calculated MW: 31.6kDa

Antigen Information

Gene Name	CASP3
Alternative Names	CASP3; Caspase 3; Apopain; CPP32; CPP32B; Caspase 3, Apoptosis-Related Cysteine Peptidase; Caspase 3, Apoptosis-Related Cysteine Protease; SREBP Cleavage Activity 1; Cysteine Protease CPP32; Protein Yama; EC 3.4.22.56; Caspase-3; CASP-3; CPP-32; SCA-1; Yama; PARP Cleavage Protease; Procaspase3; EC 3.4.22
Gene ID	836.0
SwissProt ID	P42574
Immunogen	A synthesized peptide derived from human pro Caspase 3

Background


The protein encoded by this gene is a cysteine-aspartic acid protease that plays a central role in the execution-phase of cell

apoptosis. The encoded protein cleaves and inactivates poly(ADP-ribose) polymerase while it cleaves and activates sterol regulatory element binding proteins as well as caspases 6, 7, and 9. This protein itself is processed by caspases 8, 9, and 10. It is the predominant caspase involved in the cleavage of amyloid-beta 4A precursor protein, which is associated with neuronal death in Alzheimer's disease. [provided by RefSeq, Aug 2017]


Research Area

Cell Biology,Cancer,Metabolism,Neuroscience


Image Data




Western blotting analysis using pro Caspase 3 antibody (KVA00230). pro Caspase 3 expression in wild type (WT) and pro Caspase 3 shRNA knockdown (KD) HAP1 cells with 20 µg of total cell lysates. β-Tubulin serves as a loading control. The blot was incubated with pro Caspase 3 antibody (KVA00230, 1:5,000) and HRP-conjugated goat anti-rabbit secondary antibody (APS0635, 1:10,000) respectively.




Western blotting analysis using pro Caspase 3 antibody (KVA00230). Total cell lysates (30 µg) from various cell lines were loaded and separated by SDS-PAGE. The blot was incubated with pro Caspase 3 antibody (KVA00230, 1:5,000) and HRP-conjugated goat anti-rabbit secondary antibody (APS0635, 1:10,000) respectively.



Immunohistochemistry was performed on paraffin-embedded human tonsillitis using pro caspase 3 antibody (KVA00230, 1:200). Antigen retrieval was done in sodium citrate buffer (pH 6.0). DAB was used for detection, with hematoxylin counterstaining. Images were acquired using a Nikon Ci-L Plus microscope (40× objective). Scale bar: 25 µm.



Validation of pro Caspase 3 knockdown using flow cytometry. Wild-type(WT, Blue) and knockdown(KD, Green) HeLa cells were stained with pro Caspase 3 antibody (KVA00230, 1:2,000) and analyzed using CytoFLEX.



Flow cytometric analysis of pro Caspase 3 expression in HAP-1 cells using pro Caspase 3 antibody (KVA00230, 1:2,000). Green, isotype control; red, pro Caspase 3.