

Product Name: KD-Validated PRAS40 Recombinant Rabbit Monoclonal Antibody
Catalog #: KVA00240

For research use only.

Summary

Description	KO&KD-Validated antibody
Host	Rabbit
Application	WB,FCM,ICC
Reactivity	Human,Mouse,Rat
Conjugation	Unconjugated
Modification	Unmodified
Isotype	Rabbit IgG
Clonality	Rabbit mAb
Form	Liquid
Storage	Aliquot and store at -20°C (valid for 12 months). Avoid freeze/thaw cycles.
Shipping	Ice bags
Buffer	Supplied in PBS (pH 7.4) containing 50% glycerol, and 0.02% sodium azide.
Purification	Affinity purification

Application

Dilution Ratio	WB 1:1,000-1:2,500; FC 1:200-1:2,000; ICC 1:100-1:1,000
Molecular Weight	Calculated MW: 27.4kDa

Antigen Information

Gene Name	AKT1S1 AKT1 Substrate 1; PRAS40; Lobe; 40 KDa Proline-Rich AKT Substrate; Proline-Rich AKT1
Alternative Names	Substrate 1; MGC2865; Proline-Rich Akt Substrate, 40 KDa; AKT1 Substrate 1 (Proline-Rich); AKT1 Substrate 1 (Proline Rich)
Gene ID	84335.0
SwissProt ID	Q96B36
Immunogen	A synthesized peptide derived from human PRAS40


Background

AKT1S1 is a proline-rich substrate of AKT (MIM 164730) that binds 14-3-3 protein (see YWHAH, MIM 113508) when phosphorylated (Kovacina et al., 2003 [PubMed 12524439]).[supplied by OMIM, Mar 2008]


Research Area

Cell Biology


Image Data




Western blotting analysis using PRAS40 antibody (KVA00240). PRAS40 expression in wild type (WT) and PRAS40 shRNA knockdown (KD) HeLa cells with 30 µg of total cell lysates. Hsp90 α serves as a loading control. The blot was incubated with PRAS40 antibody (KVA00240, 1:2,500) and HRP-conjugated goat anti-rabbit secondary antibody (APS0635, 1:10,000) respectively.



Western blotting analysis using PRAS40 antibody (KVA00240). Total cell lysates (30 µg) from various cell lines were loaded and separated by SDS-PAGE. The blot was incubated with PRAS40 antibody (KVA00240, 1:2,500) and HRP-conjugated goat anti-rabbit secondary antibody (APS0635, 1:10,000) respectively.



Immunocytochemical staining of C2C12 cells with PRAS40 antibody (KVA00240, 1:1,000). Nuclei were stained blue with DAPI; PRAS40 was stained magenta with Alexa Fluor® 647. Images were taken using Leica stellaris 5. Protein abundance based on laser Intensity and smart gain: Medium. Scale bar: 20 µm.



Flow cytometric analysis of PRAS40 expression in C2C12 cells using PRAS40 antibody (KVA00240, 1:2,000). Green, isotype control; red, PRAS40.