

Product Name: KD-Validated Cleaved-Caspase 8 Rabbit Polyclonal Antibody**Catalog #: KVA00442**

For research use only.

Summary

Description	KO&KD-Validated antibody
Host	Rabbit
Application	WB
Reactivity	Human
Conjugation	Unconjugated
Modification	Unmodified
Isotype	Rabbit IgG
Clonality	Rabbit pAb
Form	Liquid
Storage	Aliquot and store at -20°C (valid for 12 months). Avoid freeze/thaw cycles.
Shipping	Ice bags
Buffer	Supplied in PBS (pH 7.4) containing 50% glycerol, and 0.02% sodium azide.
Purification	Affinity purification

Application

Dilution Ratio	WB 1:500-1:2,500
Molecular Weight	Calculated MW: 55.4kDa

Antigen Information

Gene Name	CASP8
Alternative Names	CASP8; Caspase 8; FLICE; MCH5; MACH; Casp-8; Caspase 8, Apoptosis-Related Cysteine Peptidase; Caspase 8, Apoptosis-Related Cysteine Protease; MORT1-Associated Ced-3 Homolog; ICE-Like Apoptotic Protease 5; Apoptotic Cysteine Protease; Apoptotic Protease Mch-5; FADD-Like ICE; Caspase-8; CAP4; FADD-Homologous ICE/CED-3-Like Protease; FADD-Homologous ICE/Ced-3-Like Protease; MACH-Beta-1/2/3/4 Protein; MACH-Alpha-1/2/3 Protein; EC 3.4.22.61; ALPS2B; CASP-8
Gene ID	841.0
SwissProt ID	Q14790
Immunogen	A synthesized peptide derived from human Cleaved Caspase 8


Background

This gene encodes a member of the cysteine-aspartic acid protease (caspase) family. Sequential activation of caspases plays a central role in the execution-phase of cell apoptosis. Caspases exist as inactive proenzymes composed of a prodomain, a large protease subunit, and a small protease subunit. Activation of caspases requires proteolytic processing at conserved internal aspartic residues to generate a heterodimeric enzyme consisting of the large and small subunits. This protein is involved in the programmed cell death induced by Fas and various apoptotic stimuli. The N-terminal FADD-like death effector domain of this protein suggests that it may interact with Fas-interacting protein FADD. This protein was detected in the insoluble fraction of the affected brain region from Huntington disease patients but not in those from normal controls, which implicated the role in neurodegenerative diseases. Many alternatively spliced transcript variants encoding different isoforms have been described, although not all variants have had their full-length sequences determined. [provided by RefSeq, Jul 2008]

Research Area

Cell Biology,Cancer,Metabolism

Image Data



Western blotting analysis using caspase 8 antibody (KVA00442). Total cell lysates (30 µg) from various cell lines were loaded and separated by SDS-PAGE. The blot was incubated with caspase 8 antibody (KVA00442, 1:2,500) and HRP-conjugated goat anti-rabbit secondary antibody (APS0635, 1:10,000) respectively.

