

Product Name: KD-Validated Beclin 1 Recombinant Rabbit Monoclonal Antibody
Catalog #: KVA00772

For research use only.

Summary

Description	KO&KD-Validated antibody
Host	Rabbit
Application	WB,FCM
Reactivity	Human,Mouse,Rat
Conjugation	Unconjugated
Modification	Unmodified
Isotype	Rabbit IgG
Clonality	Rabbit mAb
Form	Liquid
Storage	Aliquot and store at -20°C (valid for 12 months). Avoid freeze/thaw cycles.
Shipping	Ice bags
Buffer	Supplied in PBS (pH 7.4) containing 50% glycerol, and 0.02% sodium azide.
Purification	Affinity purification

Application

Dilution Ratio	WB 1:1,000-1:5,000; FC 1:200-1:2,000
Molecular Weight	Calculated MW: 51.9kDa

Antigen Information

Gene Name	BECN1 BECN1; Beclin 1; VPS30; ATG6; Coiled-Coil Myosin-Like BCL2-Interacting Protein; Beclin 1, Autophagy Related; Beclin-1; Beclin 1 (Coiled-Coil, Moesin-Like BCL2 Interacting Protein);
Alternative Names	Beclin 1 (Coiled-Coil, Moesin-Like BCL2-Interacting Protein); ATG6 Autophagy Related 6 Homolog (S. Cerevisiae); Testis Secretory Sperm-Binding Protein Li 215e; ATG6 Autophagy Related 6 Homolog; Beclin1-Gamma; Protein GT197; Beclin1-Beta; Beclin1; GT197
Gene ID	8678.0
SwissProt ID	Q14457
Immunogen	Recombinant protein of human Beclin 1


Background

This gene encodes a protein that regulates autophagy, a catabolic process of degradation induced by starvation. The encoded protein is a component of the phosphatidylinositol-3-kinase (PI3K) complex which mediates vesicle-trafficking processes. This protein is thought to play a role in multiple cellular processes, including tumorigenesis, neurodegeneration and apoptosis. Alternative splicing results in multiple transcript variants. [provided by RefSeq, Sep 2015]


Research Area

Epigenetics and Nuclear Signaling, Cancer, Microbiology, Cancer, Metabolism, Neuroscience


Image Data



Western blotting analysis using beclin 1 antibody (KVA00772). Beclin 1 expression in wild type (WT) and beclin 1 (BECN1) shRNA knockdown (KD) HeLa cells with 20 µg of total cell lysates. Hsp90 α serves as a loading control. The blot was incubated with beclin 1 antibody (KVA00772, 1:5,000) and HRP-conjugated goat anti-rabbit secondary antibody (APS0635, 1:10,000) respectively.



Western blotting analysis using beclin 1 antibody (KVA00772). Total cell lysates (30 µg) from various cell lines were loaded and separated by SDS-PAGE. The blot was incubated with beclin 1 antibody (KVA00772, 1:5,000) and HRP-conjugated goat anti-rabbit secondary antibody (APS0635, 1:10,000) respectively.



Validation of Beclin 1 knockdown using flow cytometry. Wild-type (WT, Blue) and knockdown (KD, Green) HeLa cells were stained with Beclin 1 antibody (KVA00772, 1:2,000) and analyzed using BD flow cytometer.