

Product Name: KD-Validated Cyclin B1 Recombinant Rabbit Monoclonal Antibody
Catalog #: KVAb00796

For research use only.

Summary

Description	KO&KD-Validated antibody
Host	Rabbit
Application	WB,FCM,ICC
Reactivity	Human,Mouse,Rat
Conjugation	Unconjugated
Modification	Unmodified
Isotype	Rabbit IgG
Clonality	Rabbit mAb
Form	Liquid
Storage	Aliquot and store at -20°C (valid for 12 months). Avoid freeze/thaw cycles.
Shipping	Ice bags
Buffer	Supplied in PBS (pH 7.4) containing 50% glycerol, and 0.02% sodium azide.
Purification	Affinity purification

Application

Dilution Ratio	WB 1:1,000-1:5,000; FC 1:200-1:2,000; ICC 1:100-1:1,000
Molecular Weight	Calculated MW: 48.3kDa

Antigen Information

Gene Name	CCNB1
Alternative Names	CCNB1; Cyclin B1; CCNB; G2/Mitotic-Specific Cyclin-B1; G2/Mitotic-Specific Cyclin B1
Gene ID	891.0
SwissProt ID	P14635
Immunogen	Recombinant protein of mouse Cyclin B1


Background

The protein encoded by this gene is a regulatory protein involved in mitosis. The gene product complexes with p34(cdc2) to form the maturation-promoting factor (MPF). The encoded protein is necessary for proper control of the G2/M transition phase of the cell cycle. [provided by RefSeq, Aug 2017]


Research Area

Cell Biology, Epigenetics and Nuclear Signaling, Cancer


Image Data




Western blotting analysis using cyclin B1 antibody (KVA00796). Cyclin B1 expression in wild-type (WT) and cyclin B1 (CCNB1) shRNA knockdown (KD) HT-1080 cells with 20 µg of total cell lysates. Hsp90 α serves as a loading control. The blot was incubated with cyclin B1 antibody (KVA00796, 1:5,000) and HRP-conjugated goat anti-rabbit secondary antibody (APS0635, 1:10,000) respectively.



Western blotting analysis using cyclin B1 antibody (KVA00796). Total cell lysates (30 µg) from various cell lines were loaded and separated by SDS-PAGE. The blot was incubated with cyclin B1 antibody (KVA00796, 1:5,000) and HRP-conjugated goat anti-rabbit secondary antibody (APS0635, 1:10,000) respectively.



Immunocytochemical staining of HeLa cells with cyclin B1 antibody (KVA00796, 1:1,000). Nuclei were stained blue with DAPI; Cyclin B1 was stained magenta with Alexa Fluor® 647. Images were taken using Leica stellaris 5. Protein abundance based on laser Intensity and smart gain: High. Scale bar: 20 µm.



Flow cytometric analysis of Cyclin B1 expression in HeLa cells using Cyclin B1 antibody (KVA00796, 1:2,000). Green, isotype control; red, Cyclin B1.